



©AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2021.1D

Diagnostik und Therapie früher und fortgeschritten Mammakarzinome

Komplementäre Therapie „Survivorship“

FORSCHEN
LEHREN
HEILEN

Screened Data Sources:

Pubmed	2012 - 01/2020
ASCO	2015 – 2019
SABCS	2015 – 2019
EBCC	2015 – 2019
Cochrane library:	summary Jan. 2020:



Komplementäre Therapien

Hormontherapie „Survivorship“ (Rezidiv-Prävention)

- **Versionen 2002–2020:**


Albert / Bauerfeind / Blohmer / Fersis / Friedrich / Gerber / Göhring
/ Hanf / Janni / Kümmel / Lück / von Minckwitz / Nitz / Oberhoff /
Rhiem / Scharl / Schmidt / Schütz / Thomssen / Kümmel / Schütz

- **Version 2021:**


Dall / Solomayer

Screened Data Sources:

Pubmed	2012 - 01/202
ASCO	2015 – 2020
SABCS	2015 – 2020
EBCC	2015 – 2020
Cochrane library:	summary Jan. 2021:



CAM



Komplementäre Verfahren werden parallel zur konventionellen Therapie angewendet und unterscheiden sich von alternativen Verfahren dadurch, dass sie den Wert der konventionellen Verfahren nicht in Frage stellen, sondern sich als Ergänzung verstehen

Onkolleitlinienprogramm

1. Witt CM et al.. A Comprehensive Definition for Integrative Oncology. J Natl Cancer Inst Monogr (2017) 2017(52): Igx012

“Integrative oncology is a patient-centered, evidence-informed field of cancer care that utilizes mind and body practices, natural products, and/or lifestyle modifications from different traditions alongside conventional cancer treatments. Integrative oncology aims to optimize health, quality of life, and clinical outcomes across the cancer care continuum and to empower people to prevent cancer and become active participants before, during, and beyond cancer treatment.”



©AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2021.1D

www.ago-online.de

FORSCHEN
LEHREN
HEILEN

Gute klinische Praxis

- **Alle Patienten sollen frühestmöglich und im Verlauf wiederholt zum Interesse an Informationen komplementärmedizinischer Maßnahmen befragt werden und bei Interesse soll auf verlässliche Informationsquellen verwiesen werden.**

S3 LL "Komplementärmedizin in der Behandlung von onkologischen PatientInnen"

Literatur S3 LL "Komplementärmedizin in der Behandlung von onkologischen PatientInnen"



Allgemein

	Oxford		
	LoE	GR	AGO
■ CAM anstelle lokoregionärer Interventionen	2b	B	--
■ CAM anstelle systemischer Therapie	2b	B	--
■ Patienten sollten nach ihrer Nutzung von komplementären und alternativen Therapien befragt werden.			
■ Diagnostische Verfahren im Zusammenhang mit komplementären und alternativen Therapiekonzepten ohne Evidenz (z.B. Irisdiagnostik, Bioresonanz) sollen nicht empfohlen werden.			
■ <i>Unter Systemtherapie:</i> Besondere Beachtung gilt möglichen Medikamenteninteraktionen			

1. Saquib J, Parker BA, Natarajan L, et al. Prognosis following the use of complementary and alternative medicine in women diagnosed with breast cancer. *Complement Ther Med.* 2012 Oct;20(5):283-90. doi: 10.1016/j.ctim.2012.04.002. Epub 2012 Apr 27.
2. Guha N, Kwan ML, Quesenberry CP, et al: Soy isoflavones and risk of cancer recurrence in a cohort of breast cancer survivors: the Life After Cancer Epidemiology study. *Breast Cancer Res Treat.* 2009;118(2):395–405, pmid:19221874.
3. Johnson SB et al.. Use of Alternative Medicine for Cancer and Its Impact on Survival. *JNCI J Natl Cancer Inst* (2018) 110(1): djx145.
4. Fremd C et al.. Use of complementary and integrative medicine among German breast cancer patients: predictors and implications for patient care within the PRAEGNANT study network. *Arch Gynecol Obstet.* 2017 May;295(5):1239-1245. doi: 10.1007/s00404-017-4348-2. Epub 2017 Mar 22.
5. Samuels N et al.. Unmonitored use of herbal medicine by patients with breast cancer: reframing expectations. *J Cancer Res Clin Oncol* (2017) 143:2267–2273
6. Smith PJ et al.. Complementary and alternative medicine use by patients receiving curative-intent chemotherapy. *Asia-Pacific Journal of Clinical Oncology* 2016; 12: 265–274
7. Greenlee H et al.. Association Between Complementary and Alternative Medicine Use and Breast Cancer Chemotherapy Initiation: The Breast Cancer Quality of Care (BQUAL) Study. *JAMA Oncol.* 2016 Sep 1;2(9):1170-6. doi: 10.1001/jamaoncol.2016.0685
8. Johnson SB et al. Complementary Medicine, Refusal of Conventional Cancer Therapy, and Survival Among Patients With Curable Cancers. *JAMA Oncol.* doi:10.1001/jamaoncol.2018.2487 published online July 19, 2018.

9. Johnson SB, Park HS, Gross CP et al. Use of Alternative Medicine for Cancer and Its Impact on Survival. *J Natl Cancer Inst.* 2018 Jan 1;110(1). doi: 10.1093/jnci/djx145.
10. Hack, et al. Analysis of Oncological second opinions in a certified university breast and gynecological cancer center in relation to complementary and alternative medicine . *Medicine Research*, 2020



Komplementäre Therapien prä- und postoperativ

	Oxford	LoE	GR	AGO
Präoperativ				
▪ Hypnose (reduziert Ängste, Schmerz, Übelkeit)		1b	B	+
Postoperativ				
▪ Akupunktur				
▪ bei Schmerzen, Ängstlichkeit		1b	B	+/-
▪ bei Übelkeit, Erbrechen		2b	B	+
▪ Massage Therapie (bei Schmerzen)		2b	C	+/-
▪ Frühzeitige postoperative Bewegungstherapie beugt Dysfunktion der oberen Extremität vor CAVE: vermehrt Wundsekret		1a	A	+
▪ Körperliche Aktivität				
▪ zur Reduktion des sek. Lymphödems		1a	A	+
▪ zur Prophylaxe eines Lymphödems		1b	B	+/-
▪ Prophylaktische Lymphdrainage		1b	B	-
▪ Yoga (bei Arm- und Schulterschmerzen)		2b	C	+
▪ Musiktherapie (Schmerzreduktion nach Mastektomie)		2b	C	+/-

Hypnosis

1. Amraoui J, Pouliquen C, Fraisse J et al. (2018) Effects of a Hypnosis Session Before General Anesthesia on Postoperative Outcomes in Patients Who Underwent Minor Breast Cancer Surgery: The HYPNOSEIN Randomized Clinical Trial. *JAMA Netw Open.*;1(4):e181164. doi: 10.1001/jamanetworkopen.2018.1164.
2. Cramer H, Lauche R, Paul A, et al: Hypnosis in Breast Cancer Care: A Systematic Review of Randomized Controlled Trials. *Integr Cancer Ther.* 2015 Jan;14(1):5-15. Epub 2014 Sep 18.
3. Montgomery GH, David D, Kangas M, et al. (2014) Randomized Controlled Trial of a Cognitive-Behavioral Therapy Plus Hypnosis Intervention to Control Fatigue in Patients Undergoing Radiotherapy for Breast Cancer. *JCO DOI 10.1200/JCO.2013.49.3437*
4. Montgomery GH, Bovbjerg DH, Schnur JB et al. (2007): A randomized clinical trial of a brief hypnosis intervention to control side effects in breast surgery patients. *J Nat Cancer Inst;* 99:1304–1312.
5. Schnur JB, Bovbjerg DH, David D et al. (2008): Hypnosis decreases presurgical distress in excisional breast biopsy patients. *Anesth Analg*, 106(2):440-4
6. Montgomery GH, Schnur JB, Kravits K. Hypnosis for cancer care: Over 200 years young. *CA Cancer J Clin.* 2012 Nov 20. doi: 10.3322/caac.21165.

Acupuncture and Postoperative Nausea and Vomiting / Pain

1. Mallory MJ et al.: Acupuncture in the postoperative setting for breast cancer patients: a feasibility study. *Am J Chin Med.* 2015;43(1):45-56.
2. Chao LF et al.: The efficacy of acupoint stimulation for the management of therapy-related adverse events in patients with breast cancer: a systematic review. *Breast Cancer Res Treat* (2009) 118:255–267.
3. Quinlan-Woodward J, Gode A, Dusek JA: Assessing the Impact of Acupuncture on Pain, Nausea, Anxiety, and Coping in Women Undergoing a Mastectomy. *Oncol Nurs Forum.* 2016 Nov 1;43(6):725-732.
4. Giron PS, Haddad CA, Lopes de Almeida: Effectiveness of acupuncture in rehabilitation of physical and functional disorders of women undergoing breast cancer surgery. *Support Care Cancer.* 2016 Jun;24(6):2491-6.
5. Chiu HY, Hsieh YJ, Tsai PS. Systematic review and meta-analysis of acupuncture to reduce cancer-related pain. *Eur J Cancer Care (Engl).* 2017 Mar;26(2). doi: 10.1111/ecc.12457. Epub 2016 Feb 7

Massage Therapy

1. Lee SH, Kim JY, Yeo S et al: Meta-Analysis of Massage Therapy on Cancer Pain. *Integr Cancer Ther.* 2015 Jul;14(4):297-304.
2. Pan YQ, Yang KH, Wang YL, et al: Massage interventions and treatment-related side effects of breast cancer: a systematic review and meta-analysis. *Int J Clin Oncol.* 2014 Oct;19(5):829-41.
3. Dilaveri CA, Croghan I, Mallory MJ, et al, Massage compared with massage plus acupuncture for breast cancer patients undergoing reconstructive surgery. *J Altern Complement Med* 2020 26(7):602-609

Postoperative exercise

1. De Groef A, Van Kampen M, Dieltjens E, et al. Effectiveness of postoperative physical therapy for upper-limb impairments after breast cancer treatment: a systematic review. *Arch Phys Med Rehabil.* 2015 Jun;96(6):1140-53. doi: 10.1016/j.apmr.2015.01.006. Epub 2015 Jan 13. Review.
2. McNeely ML, Campbell K, Ospina M et al.: Exercise interventions for upper-limb dysfunction due to breast cancer treatment. *Cochrane Database of Systematic Reviews* 2010, Issue 6. Art. No.: CD005211. DOI: 10.1002/14651858.CD005211.pub2.
3. Cavanaugh KM.: Effects of Early Exercise on the Development of Lymphedema in Patients With Breast Cancer Treated With Axillary Lymph Node Dissection. *J Oncol Pract.* 2011 March; 7(2): 89–93.
4. Anderson RT, Kimmick GG, McCoy TP, et al. A randomized trial of exercise on well-being and function following breast cancer surgery: the RESTORE trial. *J Cancer Surv* 2012;6(2):172-81
5. Eyigor S, Uslu R, Apaydin S, et al. Can Yoga have any effect on shoulder and arm pain and quality of life in patients with breast cancer?

A randomized, controlled, single-blind trial . Complementary Therapies in Clinical Practice 2018;32:40-45.

Reduction secondary lymphedema

1. Baumann FT, Reike A, Reimer V et al: Effects of physical exercise on breast cancer –related secondary lymphedema : a systematic review Br Ca res Treatment 2018; 170: 1-13

Prevention lymphedema

1. Ammitzbøll G, Johansen C, Lanng C, Andersen EW et al.. Progressive resistance training to prevent arm lymphedema in the first year after breast cancer surgery: Results of a randomized controlled trial. Cancer. 2019 May 15;125(10):1683-1692. doi: 10.1002/cncr.31962. Epub 2019 Jan 11.
2. Baumann FT, Reike A, Hallek M, et al. (2018) Does Exercise have a preventive effect on secondary lymphedema in breast cancer patients following local treatment – a systemic review. Breast Care 13(5): 380–385. DOI. 10.1159/000487428

Prophylactic lymph drainage

1. Devoogdt N, Geraerts I, Van Kampen M, et al. Manual lymph drainage may not have a preventive effect on the development of breast cancer-related lymphoedema in the long term: a randomised trial. J Physiother. 2018 Oct;64(4):245-254. doi: 10.1016/j.jphys.2018.08.007. Epub 2018 Sep 18.
2. Devoogdt N, Christiaens MR, Geraerts I, et al: Effect of manual lymph drainage in addition to guidelines and exercise therapy on arm lymphoedema related to breast cancer: randomised controlled trial. BMJ 2011;343:d5326 doi: 10.1136/bmj.d5326
3. Li L, Yuan L, Chen X: Current Treatments for Breast Cancer-Related Lymphoedema: A Systematic Review. Asian Pac J Cancer Prev. 2016 Nov 1;17(11):4875-4883.

Music therapy

1. Li, X.M., Yan H, Zhou KN, et al. Effects of music therapy on pain among female breast cancer patients after radical mastectomy: results from a randomized controlled trial. Breast Cancer Res Treat, 2011. 128(2): p. 411-9.
2. Binns-Turner, P.G., Wilson LL, et al. Perioperative music and its effects on anxiety, hemodynamics, and pain in women undergoing mastectomy. Aana j, 2011. 79(4 Suppl): p. S21-7.
3. Bradt, J., et al., Music interventions for improving psychological and physical outcomes in cancer patients. Cochrane Database Syst Rev, 2016(8): p. Cd006911.



Komplementäre Therapien

Behandlungsphase – Einfluss auf Toxizität I

Bei laufender onkologischer Standardtherapie:
CAVE: Interaktionen beachten!

- **Mistellektine (Viscum album)**
zur Reduktion therapieassozierter Nebenwirkungen
- **Thymuspeptide**
verringern Risiko schwerer Infektionen
- **Ginseng**
verringert Fatigue; (Cave: interagiert mit P Enzyme, z.B. CYP3A4)
- **Ganoderma Lucidum**
verringert Fatigue; (Cave: inhibiert P Enzyme, z.B. CYP3A4)
- **L-Carnitin**
Prävention der Toxizität, verbessert periphere Neuropathie
verringert Fatigue
- **Curcumin**
vermindert Radiodermatitis
- **Ingwer**
komplementär zu Leitlinien-gerechter Medikation gegen Chemotherapie induzierte Übelkeit/Erbrechen; (Cave: Wechselwirkungen)

Oxford		
LoE	GR	AGO
1a	B	+/-
2a	B	+/-
2b	C	-
2b	C	-
1b	B	--
1b	B	-
1b	B	+/-
1b	C	+/-

General

- Li Y, Wang J, Lin F: A Methodology for Cancer Therapeutics by Systems Pharmacology-Based Analysis: A Case Study on Breast Cancer-Related Traditional Chinese Medicines. *PLoS One.* 2017 Jan 9;12(1):e0169363.
- Farahmand L, Darvishi B, Majidzadeh-A K: Naturally occurring compounds acting as potent anti-metastatic agents and their suppressing effects on Hedgehog and WNT/β-catenin signalling pathways. *Cell Prolif.* 2017 Feb;50(1). doi: 10.1111/cpr.12299.
- Cramer H, Lauche R, Klose P: Yoga for improving health-related quality of life, mental health and cancer-related symptoms in women diagnosed with breast cancer. *Cochrane Database Syst Rev.* 2017 Jan 3;1:CD010802.
- Neuhouser ML, Smith AW, George SM: Use of complementary and alternative medicine and breast cancer survival in the Health, Eating, Activity, and Lifestyle Study. *Breast Cancer Res Treat.* 2016 Dec;160(3):539-546.

Mistletoe

- Ostermann T, Appelbaum S, Poier D, et al.: A Systematic Review and Meta-Analysis on the Survival of Cancer Patients Treated with a Fermented Viscum album L. Extract (Iscador) – an Update of Findings. *Compl Med Res.* 2019. In press.
- Freudenthal M, Keinki C, Micke O, et al.: Mistletoe in oncological treatment: a systematic review : Part 1: survival and safety. *J Cancer Res Clin Oncol.* 2019 Mar;145(3):695-707
- Freudenthal M, Keinki C, Kutschner S, et al.: Mistletoe in oncological treatment: a systematic review : Part 2: quality of life and toxicity

- of cancer treatment. *J Cancer Res Clin Oncol.* 2019;145(4):927-939.
4. Loef M, Walach H. Quality of life in cancer patients treated with mistletoe: a systematic review and meta-analysis. *Compl Med Res.* 2019. In press.
 5. Weissenstein U, Kunz M, Oufir M, et al.: Absence of herb-drug interactions of mistletoe with the tamoxifen metabolite (E/Z)-endoxifen and cytochrome P450 3A4/5 and 2D6 in vitro. *BMC Complement Altern Med.* 2019;19:23.
 6. Thronicke A, Steele ML, Grah C, et al.: Clinical safety of combined therapy of immune checkpoint inhibitors and Viscum album L. therapy in patients with advanced or metastatic cancer. *BMC CAM.* 2017;17:534.
 7. Shneerson C, Taskila T, Gale N, et al: The effect of complementary and alternative medicine on the quality of life of cancer survivors: A systematic review and meta-analyses. *Complementary therapies in medicine* 2013;21:417-429.
 8. Pelzer F, Tröger W. Complementary Treatment with Mistletoe Extracts During Chemotherapy: Safety, Neutropenia, Fever, and Quality of Life assessed in a randomized study. *JAC* 2018;24:954-961.

Thymus

1. Wolf E, Milazzo S, Boehm K, et al. Thymic peptides for treatment of cancer patients. *Cochrane Database of Systematic Reviews* 2012, Issue 2. Art. No.: CD003993. DOI: 10.1002/14651858.CD003993.pub3.

Ginseng, Ganoderma lucidum

1. Leggett S1, Koczwara B, Miller M. The impact of complementary and alternative medicines on cancer symptoms, treatment side effects, quality of life, and survival in women with breast cancer--a systematic review. *Nutr Cancer.* 2015;67(3):373-91.
2. Jin X, Ruiz Beguerie J, Sze Daniel M-y et al: *Ganoderma lucidum* (reishi mushroom) for cancer treatment. *Cochrane Database of Systematic Reviews* 2012
3. Karimi N, Roshan VD: Change in adiponectin and oxidative stress after modifiable lifestyle interventions in breast cancer cases. *Asian Pacific journal of cancer prevention : APJCP* 2013;14:2845-2850.

L-Carnitine

1. Hershman DL, Unger JM, Crew K et al:. Randomized double-blind placebo-controlled trial of acetyl-L-carnitine for the prevention of taxane-induced neuropathy in women undergoing adjuvant breast cancer therapy. *J Clin Oncol.* 2013 Jul 10;31(20):2627-33
2. Cruciani RA, Zhang JJ, Manola J et al. L-carnitine supplementation for the management of fatigue in patients with cancer: an eastern

cooperative oncology group phase III, randomized, double-blind, placebo-controlled trial. J Clin Oncol. 2012 Nov 1;30(31):3864-9

Curcumin

1. Kumar P, Kadakol A, Shastrula P, et al: Curcumin as an adjuvant to breast cancer treatment. Anti-cancer agents in medicinal chemistry 2015
2. Bandyopadhyay D: Farmer to pharmacist: Curcumin as an anti-invasive and antimetastatic agent for the treatment of cancer. Frontiers in chemistry 2014;2:113.

Ingwer

1. Thamlikitkul L, Srimuninnim. Efficacy of ginger for prophylaxis of chemotherapy-induced nausea and vomiting in breast cancer patients receiving adriamycin-cyclophosphamide regimen: a randomized, double-blind, placebo-controlled, crossover study. Support Care in Cancer. 2017 Feb;25(2):459-464. doi: 10.1007/s00520-016-3423-8. Epub 2016 Oct 6.
2. Sanaati F, Najafi S, Kashaninia Z, et al. Effect of Ginger and Chamomile on Nausea and Vomiting Caused by Chemotherapy in Iranian Women with Breast Cancer. Asian Pac J Cancer Prev. 2016;17(8):4125-9.



Komplementäre Therapien

Behandlungsphase – Einfluss auf Toxizität II

Oxford		
LoE	GR	AGO
1b	B	-
2b	B	+/-
1b	C	-
2b	D	-
1b	B	-
1b	B	-
3b	B	-
1b	B	-*inf
5	D	--
3b	C	+/-*

* inf: Infusion in Deutschland nicht geprüfter Substanzen
 * Studienteilnahme empfohlen

General

1. Zhu L, Li L, Li Y: Chinese Herbal Medicine as an Adjunctive Therapy for Breast Cancer: A Systematic Review and Meta-Analysis. Evid Based Complement Alternat Med. 2016;2016:9469276. doi: 10.1155/2016/9469276.
2. McPherson L, Cochrane S, Zhu X: Current Usage of Traditional Chinese Medicine in the Management of Breast Cancer: A Practitioner's Perspective. Integr Cancer Ther. 2016 Sep;15(3):335-42. doi: 10.1177/1534735415607656.

Antioxidant supplements

1. Jung AY, Cai X, Thoene K, Obi N et al. Antioxidant supplementation and breast cancer prognosis in postmenopausal women undergoing chemotherapy and radiation therapy. Am J Clin Nutr. 2019 Jan 1;109(1):69-78. doi: 10.1093/ajcn/nqy223.
2. Harvie M: Nutritional supplements and cancer: Potential benefits and proven harms. American Society of Clinical Oncology educational book / ASCO American Society of Clinical Oncology Meeting 2014:e478-486.
3. van Dalen EC, Caron HN, Dickinson HO, et al: Cardioprotective interventions for cancer patients receiving anthracyclines. Cochrane Database Syst Rev 2011:Cd003917.
4. The Protective Role of Phenolic Compounds Against Doxorubicin-induced Cardiotoxicity: A Comprehensive Review. Razavi-Azarkhiavi K, Iranshahy M, Sahebkar A, et al. Nutr Cancer. 2016 Aug-Sep;68(6):892-917. doi: 10.1080/01635581.2016.1187280. Epub 2016 Jun 24. Review

Vitamin C

1. Heaney M, Gardner J, Karasavvas N et al.: Vitamin C antagonizes the cytotoxic effects of antineoplastic drugs. *Cancer Res.* 2008 Oct 1;68(19):8031-8.
2. PLoS One. 2015 Apr 7;10(4):e0120228. doi: 10.1371/journal.pone.0120228. eCollection 2015. High-dose intravenous vitamin C combined with cytotoxic chemotherapy in patients with advanced cancer: a phase I-II clinical trial. Hoffer LJ¹, Robitaille L², Zakarian R³, et al..
3. Sci Transl Med. 2014 Feb 5;6(222):222ra18. doi: 10.1126/scitranslmed.3007154. High-dose parenteral ascorbate enhanced chemosensitivity of ovarian cancer and reduced toxicity of chemotherapy. Ma Y¹, Chapman J, Levine M, et al.

Selen

1. Dennert G, Horneber M. Selenium for alleviating the side effects of chemotherapy, radiotherapy and surgery in cancer patients. *Cochrane Database of Systematic Reviews* 2010, Issue 11. Art. No.: CD005037. DOI: 10.1002/14651858.CD005037.pub2.

Coenzym Q10

1. Lesser GJ, Case D, Stark N, et al. A randomized, double-blind, placebo-controlled study of oral coenzyme Q10 to relieve self-reported treatment-related fatigue in newly diagnosed patients with breast cancer. *J Support Oncol* 2013;11(1):31-42
2. Ann Oncol. 2017 Mar 1;28(3):628-633. doi: 10.1093/annonc/mdw671. Interventions for preventing cardiomyopathy due to anthracyclines: a Bayesian network meta-analysis. Abdel-Qadir H^{1,2,3,4}, Ong G⁴, Fazelzad R⁵, Amir E^{1,5,6}, Lee DS^{1,2,6,7}, Thavendiranathan P^{6,7}, Tomlinson G^{1,6}.

Proteolytic enzymes and toxicity of chemotherapy

1. Petru U, Stranz B, Petru C: Effects of proteolytic enzyme therapy with Wobe Mugs against chemotherapy-induced toxicity in breast cancer patients - results of a pilot study. *Wien Med Wochenschr.* 2010 Nov;160(19-20):513-6.

Bromelain

1. Hidaka M, Nagata M, Kawano Y, et al.: Inhibitory effects of fruit juices on cytochrome P450 2C9 activity in vitro. *Biosci Biotechnol Biochem.* Feb 2008;72(2):406-411.

Chinese herbal medicine and wound healing

1. Chen J, Lv Q, Yu M et al.: Randomized clinical trial of Chinese herbal medications to reduce wound complications after mastectomy for breast carcinoma. Br J Surg. 2010 Dec;97(12):1798-804

Kurzzeit-Fasten

1. Bauersfeld SP, Kessler CS, Wischnewsky M et al. The effects of short-term fasting on quality of life and tolerance to chemotherapy in patients with breast and ovarian cancer: a randomized cross-over pilot study. BMC Cancer (2018) 18:476
2. Groot de S, Vreeswijk MPG, et al. the effects of short-term fasting on tolerance to (neo) adjuvant chemotherapy in Her2-negative breast cancer patients: a randomized pilot study. BMC Cancer 2015;15:652



Komplementäre Therapien unter onkologischer Therapie Behandlung von Nebenwirkungen

	Oxford		
	LoE	GR	AGO
▪ Chinesische Kräutermedizin (zur Behandlung chemo-therapiebedingter Nebenwirkungen)	1b	B	-
▪ Homöopathische Medizin (gegen therapiebedingte Nebenwirkungen) (Placeboeffekt)	1b	B	+/-
▪ Topische Anwendung Silymarin (Silybin, Mariendistelextrakt) (akute Hautreaktion unter Strahlentherapie)	3a	B	+/-
▪ Massage (zur Verbesserung von Fatigue, Schmerzen, Angst, Übelkeit)	1b	C	+/-
▪ Transkutane elektrische Nervenstimulation (TENS) (bei Karzinomschmerzen)	2b	D	+/-
▪ Hydrotherapie (bei therapie-assozierter Hauttrockenheit, Hautverhärtung, Nagelveränderung)	3b	C	+/-

Chinese medicinal herbs

1. Zhu L, Li L, Li Y: Chinese Herbal Medicine as an Adjunctive Therapy for Breast Cancer: A Systematic Review and Meta-Analysis. Evid Based Complement Alternat Med. 2016;2016:9469276. doi: 10.1155/2016/9469276.

Homeopathic medicines for adverse effects of cancer treatments

1. Kassab S, Cummings M, Berkovitz S, et al. Homeopathic medicines for adverse effects of cancer treatments. Cochrane Database of Systematic Reviews 2012, Issue 8. Art. No.: CD004845. DOI: 10.1002/14651858.CD004845.pub2.

Topical use of Silymarin

1. Lazzeroni M, Guerrieri-Gonzaga A, Gandini S: A Presurgical Study of Oral Silybin-Phosphatidylcholine in Patients with Early Breast Cancer. Cancer Prev Res (Phila). 2016 Jan;9(1):89-95. doi: 10.1158/1940-6207.

Massage

1. Izgu N, Metin ZG, Karadas C et al.. Prevention of chemotherapy-induced peripheral neuropathy with classical massage in breast cancer patients receiving paclitaxel: An assessor-blinded randomized controlled trial. Eur J Oncol Nurs. 2019 Jun;40:36-43. doi: 10.1016/j.ejon.2019.03.002. Epub 2019 Mar 22.

2. Shin ES, Seo KH, Lee SH, et al. Massage with or without aromatherapy for symptom relief in people with cancer. Cochrane Database of Systematic Reviews 2016, Issue 6. Art. No.: CD009873. DOI: 10.1002/14651858.CD009873.pub3.
3. Robison JG, Smith CL. Therapeutic Massage During Chemotherapy and/or Biotherapy Infusions: Patient Perceptions of Pain, Fatigue, Nausea, Anxiety, and Satisfaction. Clin J Oncol Nurs. 2016 Apr;20(2):E34-40. doi: 10.1188/16.CJON.E34-E40.
4. Donoyama N, Satoh T, Hamano T et al., Effects of Anma therapy (Japanese massage) on health-related quality of life in gynecologic cancer survivors: a randomized controlled trial. PLoS one 2018;13:e0196638.

Transcutaneous electric nerve stimulation (TENS) for cancer pain in adults: (von Slide 16 übertragen)

1. Hurlow A, Bennett MI, Robb KA, et al. Transcutaneous electric nerve stimulation (TENS) for cancer pain in adults. Cochrane Database of Systematic Reviews 2012, Issue 3. Art. No.: CD006276. DOI: 10.1002/14651858.CD006276.pub3.
2. Paley CA, Johnson MI, Tashani O et al. Acupuncture for cancer pain in adults. Cochrane Database of Systematic Reviews 2011, Issue 1. Art. No.: CD007753. DOI: 10.1002/14651858.CD007753.pub2.

Hydrotherapie

1. Dalenc F, Ribet V, Rossi AB, et al. Efficacy of a global supportive skin care programme with hydrotherapy after non-metastatic breast cancer treatment: a randomized, controlled study. Eur J Cancer Care 2018;27:doi:10.1111/ecc



Komplementäre Therapien unter onkologischer Therapie Behandlung von Nebenwirkungen

	Oxford	LoE	GR	AGO
Akupunktur zur Verbesserung von:				
▪ Chemotherapie-induzierter Übelkeit und Erbrechen		1b	B	+
▪ Elektro-Akupunktur als Ergänzung zu antiemetischer Therapie		1b	B	+
▪ Akupressur als Ergänzung zu Antiemetika				
▪ Schmerzen		1b	B	+
▪ Krebs schmerzen		1a	B	+
▪ AI-induzierter Arthralgie		2b	D	+/-
▪ TENS - transkutane elektrische Nervenstimulation bei Krebs schmerzen		1a	B	+
▪ Fatigue		1b	B	+
▪ Akupressur		2b	B	+
▪ Angst und Depression		5	D	+/-
▪ Kognitiver Dysfunktion		1b	B	+
▪ Menopausensyndrom bei Patientinnen mit Mammakarzinom		1b	B	+/-
▪ zur Verbesserung v. Häufigkeit und Schwere d. Hitzewallungen		2a	B	+
▪ Elektroakupunktur zur Verbesserung des Schlafs bei Hitzewallungen		2b	B	+/-
▪ Leukopenie (Moxibustion)		1b	B	-
▪ Chemotherapie-induzierter Polyneuropathie		2b	B	+/-
▪ als Prophylaxe		2b	B	+/-
▪ als Therapie		2b	B	+/-
▪ Chronischem Lymphödem nach MaCa Therapie				

Acupuncture

- Brinkhaus B, Kirschbaum B, Stockigt B, et al.: Prophylactic acupuncture treatment during chemotherapy with breast cancer: a randomized pragmatic trial with a retrospective nested qualitative study. *Breast cancer research and treatment* 2019, 178(3):617-628.
- Wu X, Chung VCh, Hui EP, et al: Effectiveness of acupuncture and related therapies for palliative care of cancer: overview of systematic reviews. *Sci Rep.* 2015 Nov 26;5:16776.
- Tao WW, Jiang H, Tao XM, et al. Effects of Acupuncture, Tuina, Tai Chi, Qigong, and Traditional Chinese Medicine Five-Element Music Therapy on Symptom Management and Quality of Life for Cancer Patients: A Meta-Analysis. *J Pain Symptom Manage.* 2016 Apr;51(4):728-47. doi: 10.1016/j.jpainsymman.2015.11.027. Epub 2016 Feb 12. Review.
- Lau CH, Wu X, Chung VC, et al. Acupuncture and Related Therapies for Symptom Management in Palliative Cancer Care: Systematic Review and Meta-Analysis. *Medicine (Baltimore)*. 2016 Mar;95(9):e2901. doi: 10.1097/MD.0000000000002901.

Menopausensyndrom

- Salehi A, Marzban M, Zadeh AR: Acupuncture for treating hot flashes in breast cancer patients: an updated meta-analysis. *Support Care Cancer.* 2016 Dec;24(12):4895-4899.
- Lesi G, Razzini G, Musti MA: Acupuncture As an Integrative Approach for the Treatment of Hot Flashes in Women With Breast

- Cancer: A Prospective Multicenter Randomized Controlled Trial (AcCliMaT). *J Clin Oncol.* 2016 May 20;34(15):1795-802. doi: 10.1200/JCO.2015.63.2893.
3. Chiu HY, Shyu YK, Chang PC: Effects of Acupuncture on Menopause-Related Symptoms in Breast Cancer Survivors: A Meta-analysis of Randomized Controlled Trials. *Cancer Nurs.* 2016 May-Jun;39(3):228-37. doi: 10.1097/NCC.0000000000000278.
 4. Tao WW, Jiang H, Tao XM: Effects of Acupuncture, Tuina, Tai Chi, Qigong, and Traditional Chinese Medicine Five-Element Music Therapy on Symptom Management and Quality of Life for Cancer Patients: A Meta-Analysis. *J Pain Symptom Manage.* 2016 Apr;51(4):728-47. doi: 10.1016/j.jpainsympman.2015.11.027.
 5. Chien TJ, Hsu CH, Liu CY, et al. Effect of acupuncture on hot flush and menopause symptoms in breast cancer- A systematic review and meta-analysis. *PLoS One.* 2017 Aug 22;12(8):e0180918. doi: 10.1371/journal.pone.0180918. eCollection 2017. Review.
 6. Tao WW, Tao XM, Song CL. Effects of non-pharmacological supportive care for hot flushes in breast cancer: a meta-analysis. *Support Care Cancer.* 2017 Jul;25(7):2335-2347. doi: 10.1007/s00520-017-3691-y. Epub 2017 Apr 11.

Chemotherapy-induced Nausea and Vomiting

1. Fonnebo V et al.: Acupuncture and acupressure in the treatment of chemotherapy-associated nausea and vomiting. www.cancer.org, Updated May 21, 2009.
2. Beith JM, Oh B, Chat eld MD, et al. Electroacupuncture for nausea, vomiting, and myelosuppression in women receiving adjuvant chemo- therapy for early breast cancer: a randomized controlled pilot trial. *Medical Acupuncture.* 2012;24:241–248.
3. Eghbali M et al.. The effect of auricular acupressure on nausea and vomiting caused by chemotherapy among breast cancer patients. *Complement Ther Clin Pract.* 2016 Aug;24:189-94. doi: 10.1016/j.ctcp.2016.06.006. Epub 2016 Jul 5.

Insomnia

1. Garland SN, Xie SX, DuHamel K, et al.: Acupuncture Versus Cognitive Behavioral Therapy for Insomnia in Cancer Survivors: A Randomized Clinical Trial. *Journal of the National Cancer Institute* 2019.
2. Choi TY, Kim JI, Lim HJ, Lee MS: Acupuncture for Managing Cancer-Related Insomnia: A Systematic Review of Randomized Clinical Trials. *Integrative cancer therapies* 2017, 16(2):135-146.
3. Garland SN, Xie SX, Li Q, et al.:Comparative effectiveness of electro-acupuncture versus gabapentin for sleep disturbances in breast cancer survivors with hot flashes: a randomized trial. *Menopause (New York, NY)* 2017, 24(5):517-523.

Cognitive dysfunction

1. Tong T, Pei C, Chen J et al. Efficacy of Acupuncture Therapy for Chemotherapy-Related Cognitive Impairment in Breast Cancer Patients. *Med Sci Monit.* 2018 May 8 [revised 2018 Jan 1];24:2919-2927. doi: 10.12659/MSM.909712.

Fatigue

1. Ling WM, Lui LY, So WK, et al: Effects of acupuncture and acupressure on cancer-related fatigue: a systematic review. *Oncol Nurs Forum.* 2014 Nov 1;41(6):581-92.
2. Mao JJ, Farrar JT, Bruner D et al. Electroacupuncture for fatigue, sleep, and psychological distress in breast cancer patients with aromatase inhibitor-related arthralgia: a randomized trial. *Cancer.* 2014;120(23):3744-51.
3. Smith C, Carmady B, Thornton C et al. The effect of acupuncture on post-cancer fatigue and well-being for women recovering from breast cancer: a pilot randomised controlled trial. *Complement Ther Clin Pract.* 2013;19(1):32-5.
4. Zhang Y, Lin L, Li H, et al. Effects of acupuncture on cancer-related fatigue: a meta-analysis. *Support Care Cancer.* 2017 Nov 11. doi: 10.1007/s00520-017-3955-6.
5. Zick SM, Sen A, Wyatt GK, et al. Investigation of 2 Types of Self-administered Acupressure for Persistent Cancer-Related Fatigue in Breast Cancer Survivors: A Randomized Clinical Trial. *JAMA Oncol.* 2016 Nov 1;2(11):1470-1476. doi: 10.1001/jamaoncol.2016.1867.
6. Zhang B, Dong JN, Sun P, et al. Effect of therapeutic care for treating fatigue in patients with breast cancer receiving chemotherapy. *Medicine (Baltimore).* 2017 Aug;96(33):e7750. doi: 10.1097/MD.0000000000007750.

Pain

1. He Y, Guo X, May BH et al. Clinical Evidence for Association of Acupuncture and Acupressure With Improved Cancer Pain: A Systematic Review and Meta-Analysis. *JAMA Oncol.* 2019 Dec 19. doi: 10.1001/jamaoncol.2019.5233.
2. Pan Y, Yang K, Shi X, Liang H et al.: Clinical Benefits of Acupuncture for the Reduction of Hormone Therapy-Related Side Effects in Breast Cancer Patients: A Systematic Review. *Integrative cancer therapies* 2018, 17(3):602-618.
3. Hershman DL, Unger JM, Greenlee H, et al.: Effect of Acupuncture vs Sham Acupuncture or Waitlist Control on Joint Pain Related to Aromatase Inhibitors Among Women With Early-Stage Breast Cancer: A Randomized Clinical Trial. *Jama* 2018, 320(2):167-176.
4. Garcia MK, McQuade J, Haddad R, et al. Systematic review of acupuncture in cancer care: a synthesis of the evidence. *J Clin Oncol.* 2013 Mar 1;31(7):952-60. doi: 10.1200/JCO.2012.43.5818. Epub 2013 Jan 22. (von Slide 16 übertragen)
5. Bae K, Yoo HS, Lamoury G, et al. Acupuncture for Aromatase Inhibitor-Induced Arthralgia: A Systematic Review. *Integr Cancer Ther.*

- 2015 Nov;14(6):496-502. doi: 10.1177/1534735415596573. Epub 2015 Jul 28. (von Slide 16 übertragen)
6. Paley CA1, Johnson MI, Tashani OA, et al. Acupuncture for cancer pain in adults. Cochrane Database Syst Rev. 2015 Oct 15;10:CD007753. doi: 10.1002/14651858.CD007753.pub3. (von Slide 16 übertragen)
 7. Chen L, Lin CC, Huang TW, et al. Effect of acupuncture on aromatase inhibitor-induced arthralgia in patients with breast cancer: A meta-analysis of randomized controlled trials. Breast. 2017 Jun;33:132-138. doi: 10.1016/j.breast.2017.03.015. Epub 2017 Apr 4.
 8. Chien TJ1, Liu CY, Chang YF, et al. Acupuncture for treating aromatase inhibitor-related arthralgia in breast cancer: a systematic review and meta-analysis. J Altern Complement Med. 2015 May;21(5):251-60. doi: 10.1089/acm.2014.0083. Epub 2015 Apr 27.
 9. Yang GS1, Kim HJ, Griffith K et al. Interventions for the Treatment of Aromatase Inhibitor-Associated Arthralgia in Breast Cancer Survivors: A Systematic Review and Meta-analysis. Cancer Nurs. 2017 Jul/Aug;40(4):E26-E41. doi: 10.1097/NCC.0000000000000409
 10. Chiu HY, Hsieh YJ, Tsai PS. Systematic review and meta-analysis of acupuncture to reduce cancer-related pain. Eur J Cancer Care (Engl). 2017 Mar;26(2). doi: 10.1111/ecc.12457. Epub 2016 Feb 7.

Leucopenia

1. Choi TY, Lee MS, Ernst E: Moxibustion for the treatment of chemotherapy-induced leukopenia: a systematic review of randomized clinical trials. Support Care Cancer. 2015 Jun;23(6):1819-26.

Chemotherapy induced peripheral neuropathy

1. Molassiotis A, Suen LKP, Cheng HL et al. A Randomized Assessor-Blinded Wait-List-Controlled Trial to Assess the Effectiveness of Acupuncture in the Management of Chemotherapy-Induced Peripheral Neuropathy. Integr Cancer Ther. 2019 Jan-Dec;18:1534735419836501. doi: 10.1177/1534735419836501.
2. Lu W, Giobbie-Hurder A, Freedman RA et al. Acupuncture for Chemotherapy-Induced Peripheral Neuropathy in Breast Cancer Survivors: A Randomized Controlled Pilot Trial. Oncologist. 2019 Oct 14. pii: theoncologist.2019-0489. doi: 10.1634/theoncologist.2019-0489.
3. Greenlee H, Crew KD, Capodice J et al. Randomized sham-controlled pilot trial of weekly electro-acupuncture for the prevention of taxane-induced peripheral neuropathy in women with early stage breast cancer. Breast Cancer Res Treat. 2016 Apr;156(3):453-464. doi: 10.1007/s10549-016-3759-2. Epub 2016 Mar 25.
4. Ben-Horin I, Kahan P, Ryvo L et al. Acupuncture and Reflexology for Chemotherapy-Induced Peripheral Neuropathy in Breast Cancer. Integr Cancer Ther. 2017 Sep;16(3):258-262. doi: 10.1177/1534735417690254. Epub 2017 Feb 2. (Haben wir die?)
5. Brami C, Bao T, Deng G Natural products and complementary therapies for chemotherapy-induced peripheral neuropathy: A

- systematic review. Crit Rev Oncol Hematol. 2016 Feb;98:325-34. doi: 10.1016/j.critrevonc.2015.11.014. Epub 2015 Nov 23.
6. Weidong Lu, Anita Giobbie-Hurder, Rachel A. Freedman et al, Acupuncture for Chemotherapy-Induced Peripheral Neuropathy in Breast Cancer Survivors: A Randomized Controlled Pilot Trial, Oncologist. 2020 Apr; 25(4): 310–318.

Chronisches Lymphödem

1. Chien TJ, Liu CY, Fang CJ: The Effect of Acupuncture in Breast Cancer-Related Lymphoedema (BCRL): A Systematic Review and Meta-Analysis. Integrative Cancer Therapies 2019, 18:1534735419866910.
2. Yao C1, Xu Y1, Chen L1, Jiang H1, Ki CS2, Byun JS3, Bian W1. Curr Oncol. 2016 Feb;23(1):e27-34. doi: 10.3747/co.23.2788. Epub 2016 Feb 18. Effects of warm acupuncture on breast cancer-related chronic lymphedema: a randomized controlled trial.
3. Zayas J, Ruddy KJ, Olson JE et al, Real-world experiences with acupuncture among breast cancer survivors: a cross-sectional survey study. Support Care Cancer 2020 28(12)5833-5838

Angst und Depression

1. Molassiotis, A., et al., Acupuncture for cancer-related fatigue in patients with breast cancer: a pragmatic randomized controlled trial. J Clin Oncol, 2012. 30(36): p. 4470-6.



Komplementäre Therapien Behandlungsphase – Mind-Body Medizin I

Oxford		
LoE	GR	AGO
1a	A	+

MBSR (Mindfulness-Based Stress Reduction – dt. Achtsamkeitsbasierte Stressbewältigung)

Programm verbessert Lebensqualität, Bewältigungsstrategien, Achtsamkeit, vermindert Stress, Angst, Depression, Fatigue und Schlafstörung

Körperliches Training/Sport (mind. 3x/Woche moderates

Ausdauertraining in Kombination mit kräftigendem Geräterunning 2 x /Wo.) verbessert Lebensqualität, kardiorespiratorische Fitness, körperliche Leistungsfähigkeit, Schlaf, Schmerz, Depression, Lymphödem und Fatigue

1a A ++

Mind-Body Medicine (MBM)

- Charalambous A, Giannakopoulou M, Bozas E, et al: Guided Imagery And Progressive Muscle Relaxation as a Cluster of Symptoms Management Intervention in Patients Receiving Chemotherapy: A Randomized Control Trial. PLoS One. 2016 Jun 24;11(6):e0156911. doi: 10.1371/journal.pone.0156911. eCollection 2016.

MBSR

- Schell LK, Monsef I, Wöckel A, Skoetz N. Mindfulness-based stress reduction for women diagnosed with breast cancer. Cochrane Database Syst Rev. 2019 Mar 27;3:CD011518.
- Zhang Q, Zhao H, Zheng Y. (2019) Effectiveness of mindfulness-based stress reduction (MBSR) on symptom variables and health-related quality of life in breast cancer patients-a systematic review and meta-analysis. Support Care Cancer;27(3):771-781. doi: 10.1007/s00520-018-4570-x. Epub 2018 Nov 28.
- Haller H, Winkler MM, Klose P et al: Mindfulness-based interventions for women with breast cancer: an updated systematic review and meta-analysis. Acta oncologica (Stockholm, Sweden). 2017 Jul 07:1-12. PubMed PMID: 28686520. Epub 2017/07/08. eng.
- Lengacher CA, Reich RR, Paterson CL, et al: Examination of Broad Symptom Improvement Resulting From Mindfulness-Based Stress Reduction in Breast Cancer Survivors: A Randomized Controlled Trial. J Clin Oncol. 2016 Aug 20;34(24):2827-34. doi: 10.1200/JCO.2015.65.7874. Epub 2016 May 31.

5. Huang HP, He M, Wang HY, et al. A meta-analysis of the benefits of mindfulness-based stress reduction (MBSR) on psychological function among breast cancer (BC) survivors. *Breast Cancer*. 2015 Mar 28. [Epub ahead of print]
6. Lengacher CA, Reich RR, Paterson CL: Examination of Broad Symptom Improvement Resulting From Mindfulness-Based Stress Reduction in Breast Cancer Survivors: A Randomized Controlled Trial. *J Clin Oncol*. 2016 Aug 20;34(24):2827-34. doi: 10.1200/JCO.2015.65.7874.
7. Huang HP, He M, Wang HY: A meta-analysis of the benefits of mindfulness-based stress reduction (MBSR) on psychological function among breast cancer (BC) survivors. *Breast Cancer*. 2016 Jul;23(4):568-76. doi: 10.1007/s12282-015-0604-0.
8. Reich RR, Lengacher CA, Alinat CB: Mindfulness-Based Stress Reduction in Post-treatment Breast Cancer Patients: Immediate and Sustained Effects Across Multiple Symptom Clusters. *J Pain Symptom Manage*. 2017 Jan;53(1):85-95. doi: 10.1016/j.jpainsympman.2016.08.005.
9. Schmidt ME, Wiskemann J, Ulrich CM: Self-reported physical activity behavior of breast cancer survivors during and after adjuvant therapy: 12 months follow-up of two randomized exercise intervention trials. *Acta Oncol*. 2017 Jan 13;1-10. doi: 10.1080/0284186X.2016.1275776.
10. Castanhel F, Liberali R. Mindfulness-Based Stress Reduction on breast cancer symptoms: systematic review and meta-analysis. *Einstein* 2018;16:1-10
11. Hall DL, Luberto CM, Philpotts et al., Mind-body interventions for fear of cancer recurrence: A systematic review and meta-analysis. *Psycho-Oncology* 2018;1-13.

Physical exercise

1. Campbell, KL, et al. (2019): Consensus Statement from International Multidisciplinary Roundtable Exercise Guidelines for Cancer Survivors. *Medicine & Science in Sports & Exercise*. 51(11):2375–2390
2. Kreutz C, Schmidt ME, Steindorf K (2019) Effects of physical and mind-body exercise on sleep problems during and after breast cancer treatment: a systematic review and meta-analysis. *Breast Cancer Res Treat*.;176(1):1-15. doi: 10.1007/s10549-019-05217-9. Epub 2019 Apr 6.
3. Furmaniak AC, Menig M, Markes MH. Exercise for women receiving adjuvant therapy for breast cancer. *Cochrane Database Syst Rev*. 2016 Sep 21;9:CD005001. [Epub ahead of print]
4. Mercier J, Savard J, Bernard P: Exercise interventions to improve sleep in cancer patients: A systematic review and meta-analysis.

- Sleep Med Rev. 2017 Dec;36:43-56. doi: 10.1016/j.smrv.2016.11.001. Epub 2016 Nov 10.
5. Steindorf K; Wiskemann J; Ulrich CM; et al: Effects of exercise on sleep problems in breast cancer patients receiving radiotherapy: a randomized clinical trial. Breast Cancer Res Treat. 2017; 162(3):489-499 (ISSN: 1573-7217)
 6. Cormie P, Pumpa K, Galvao DA et al. Is it safe and efficacious for women with lymphedema secondary to breast cancer to lift heavy weights during exercise: a randomised controlled trial. J Cancer Surviv. 2013 Sep;7(3):413-24.
 7. Forbes, C.C., et al., *Prevalence and correlates of strength exercise among breast, prostate, and colorectal cancer survivors*. Oncol Nurs Forum, 2015. 42(2): p. 118-27.

Statement on quality of life

1. Zhang, X., Li, Y., & Liu, D. (2019). Effects of exercise on the quality of life in breast cancer patients: a systematic review of randomized controlled trials. Supportive Care in Cancer, 27(1), 9-21.
2. Steindorf K, Schmidt ME, Klassen O et al. Randomized, controlled trial of resistance training in breast cancer patients receiving adjuvant radiotherapy: results on cancer-related fatigue and quality of life. Ann Oncol. 2014 Nov;25(11):2237-43.
3. Schmidt ME, Wiskemann J, Armbrust Pet al. Effects of resistance exercise on fatigue and quality of life in breast cancer patients undergoing adjuvant chemotherapy: A randomized controlled trial. Int J Cancer. 2014 Dec 6. doi: 10.1002/ijc.29383. [Epub ahead of print]
4. Zeng Y, Huang M, Cheng AS et al. Meta-analysis of the effects of exercise intervention on quality of life in breast cancer survivors. Breast Cancer. 2014 May;21(3):262-74.

Cardio respiratory Fitness / Physical Functioning

1. Gebruers, N., Camberlin, M., Theunissen, F. et al.: (2019). The effect of training interventions on physical performance, quality of life, and fatigue in patients receiving breast cancer treatment: a systematic review. Supportive Care in Cancer, 27(1), 109-122.
2. Jones LW et al. (2016) Exercise and Risk of Cardiovascular Events in Women With Nonmetastatic Breast Cancer. J Clin Oncol. 10;34(23):2743-9. doi: 10.1200/JCO.2015.65.6603. Epub 2016 May 23.
3. Casla S, López-Tarruella S, Jerez Y, et al. Supervised physical exercise improves VO₂max, quality of life, and health in early stage breast cancer patients: a randomized controlled trial. Breast Cancer Res Treat. 2015 Sep;153(2):371-82.
4. Courneya KS, Segal RJ, Mackey JR et al. Effects of exercise dose and type on sleep quality in breast cancer patients receiving chemotherapy: a multicenter randomized trial. Breast Cancer Res Treat. 2014 Apr;144(2):361-9.
5. Courneya KS, McKenzie DC, Gelmon K et al. A multicenter randomized trial of the effects of exercise dose and type on psychosocial

- distress in breast cancer patients undergoing chemotherapy. *Cancer Epidemiol Biomarkers Prev.* 2014 May;23(5):857-64.
6. Courneya KS, McKenzie DC, Mackey JR et al. Subgroup effects in a randomised trial of different types and doses of exercise during breast cancer chemotherapy. *Br J Cancer.* 2014 Oct 28;111(9):1718-25.

Fatigue

1. Gebruers, N., Camberlin, M., Theunissen, F. et al.: (2019). The effect of training interventions on physical performance, quality of life, and fatigue in patients receiving breast cancer treatment: a systematic review. *Supportive Care in Cancer,* 27(1), 109-122.
2. Mustian KM, Alfano CM, Heckler C, et al. (2017) Comparison of Pharmaceutical, Psychological, and Exercise Treatments for Cancer-Related Fatigue: A Meta-analysis. *JAMA Oncol.* 1;3(7):961-968. doi: 10.1001/jamaonc.2016.6914.
3. Hilfiker, R., Meichtry, A., Eicher, M et al. (2017). Exercise and other non-pharmaceutical interventions for cancer-related fatigue in patients during or after cancer treatment: a systematic review incorporating an indirect-comparisons meta-analysis. *Br J Sports Med, bjsports-2016.*
4. Lipsett A, Barrett S, Haruna F et al. The impact of exercise during adjuvant radiotherapy for breast cancer on fatigue and quality of life: A systematic review and meta-analysis. *Breast.* 2017 Apr;32:144-155. doi: 10.1016/j.breast.2017.02.002. Epub 2017 Feb 9.
5. Juvet LK, Thune I, Elvsaa IKØ, et al. The effect of exercise on fatigue and physical functioning in breast cancer patients during and after treatment and at 6 months follow-up: A meta-analysis. *Breast.* 2017 Jun;33:166-177. doi: 10.1016/j.breast.2017.04.003. Epub 2017 Apr 14.
6. Meneses-Echávez JF, González-Jiménez E, Ramírez-Vélez R. Effects of supervised exercise on cancer-related fatigue in breast cancer survivors: a systematic review and meta-analysis. *BMC Cancer.* 2015 Feb 21;15:77. doi: 10.1186/s12885-015-1069-4.
7. van Waart H, Stuiver MM, van Harten WH et al. Effect of Low-Intensity Physical Activity and Moderate- to High-Intensity Physical Exercise During Adjuvant Chemotherapy on Physical Fitness, Fatigue, and Chemotherapy Completion Rates: Results of the PACES Randomized Clinical Trial. *J Clin Oncol.* 2015 Jun 10;33(17):1918-27.
8. Zou LY, Yang L, He XL et al. Effects of aerobic exercise on cancer-related fatigue in breast cancer patients receiving chemotherapy: a meta-analysis. *Tumour Biol.* 2014 Jun;35(6):5659-67.

PNP

1. Steckmann F, Zopf EM, Lehmann HC, et al. (2014) Exercise Intervention studies in patients with PNP: a systemic review. *SportsMed.* DOI 10.1007/s40279-014-0207-5
2. Streckmann F, Lehmann HC, Balke M, et al., Sensorimotor training and whole-body vibration training have the potential to reduce

motor and sensory symptoms of chemotherapy-induced peripheral neuropathy – a randomized controlled pilot trial. Supp Care in Cancer doi.org/10.1007/s00520-018-4531-4

Schmerz

1. Irwin ML, Carmel B, Gross CP, et al. Randomized exercise Trial of AI-Induced Arthralgia in Breast cancer survivors- JCO 2015;33:1104-11011

Lymphödem

1. Baumann FG, Reike A, Reimer V et al., Effects of physical exercise on breast cancer-related secondary lymphedema: a systemic review. Breast Cancer Res Treat Doi 10.1007/s10549-018-4725
2. Baumann FT, Reike A, Hallek M, et al. Does Exercise have a preventive effect on secondary lymphedema in breast cancer patients following local treatment – a systemic review. Breast Care DOI. 10.1159/000487428

Körpergewicht

1. Mutschler NS, Scholz C et al. (2018) Prognostic Impact of Weight Change During Adjuvant Chemotherapy in Patients With High-Risk Early Breast Cancer: Results From the ADEBAR Study. Clin Breast Cancer. 18(2): 175–183



Komplementäre Therapien

Behandlungsphase – Mind-Body Medizin II

	Oxford		
	LoE	GR	AGO
Entspannungsverfahren Reduktion von Angst und Übelkeit, Verbesserung der Lebensqualität, Verminderung psychischer Belastung	2b	C	+/-
Yoga Verbesserung von Lebensqualität, Stress, Fatigue, Schlaf, Angst und Depression	1b	A	+
Qigong Verbesserung von Lebensqualität, Fatigue, Stimmung	2a	B	+/-
Tai-Chi Verbesserung von Lebensqualität, Muskelkraft, Schlaf	2a	B	+/-
Hypnose (in Kombination mit kognitiver Therapie) Verbesserung von Fatigue unter Radiotherapie, Reduktion von Distress	1b	A	+

General

1. Merckaert I, Lewis F, Delevallez F: Improving anxiety regulation in patients with breast cancer at the beginning of the survivorship period: A randomized clinical trial comparing the benefits of single-component and multi-component group interventions. *Psychooncology*. 2016 Oct 8. doi: 10.1002/pon.4294.
 2. Pan Y, Yang K, Wang Y, et al.: Could yoga practice improve treatment-related side effects and quality of life for women with breast cancer? A systematic review and meta-analysis. *Asia Pac J Clin Oncol*. 2015 Jan 6. doi: 10.1111/ajco.12329. [Epub ahead of print]

Entspannungsverfahren

1. Dikmen, H. A., & Terzioglu, F. (2019). Effects of Reflexology and Progressive muscle relaxation on pain, fatigue, and quality of life during chemotherapy in gynecologic cancer patients. *Pain Management Nursing*, 20(1), 47-53.
 2. Abbasi B, Mirzakhany N, Oshnari A et al. The effect of relaxation techniques on edema, anxiety and depression in post-mastectomy lymphedema patients undergoing comprehensive decongestive therapy: A Clinical Trail. *PLoS One* 2018;13:e0190231
 3. Lyman GH, Grennlee H et al. Integrative Therapies during and after Breast Cancer Treatment: ASCO Endorsement of the SIO clinical practice guideline. *JCO* 2018;36:2647-2655.
 4. Tian X, Tang R-Y, Xu LL et al. Progressive muscle relaxation is effective in preventing and alleviating of chemotherapy-induced nausea and vomiting among cancer patients: a systematic review of six randomized controlled trials. *Support Care Cancer* 2020 28(9):4051-

Yoga

1. Dong B, Xie C, Jing X, Lin L, Tian L. Yoga has a solid effect on cancer-related fatigue in patients with breast cancer: a meta-analysis. *Breast Cancer Res Treat.* 2019 Aug;177(1):5-16. doi: 10.1007/s10549-019-05278-w. Epub 2019 May 24. Review.
2. Cramer H, Lauche R, Klose P, et al. Yoga for improving health-related quality of life, mental health and cancer-related symptoms in women diagnosed with breast cancer. *The Cochrane database of systematic reviews.* 2017 Jan 03;1:CD010802. PubMed PMID: 28045199. Epub 2017/01/04. eng.
3. Chakrabarty J, Vidyasagar MS, Fernandes D, et al: Effectiveness of pranayama on cancer-related fatigue in breast cancer patients undergoing radiation therapy: A randomized controlled trial, *Int J Yoga.* 2015 Jan-Jun; 8(1): 47–53. doi: 10.4103/0973-6131.146062 PMCID: PMC4278135
4. Derry et.al. Yoga and self-reported cognitive problems in breast cancer survivors: a randomized controlled trial. *Psychooncology.* 2015 Aug;24(8):958-66.
5. Sprod LK, Fernandez ID, Janelsins MC, et al: Effects of yoga on cancer-related fatigue and global side-effect burden in older cancer survivors. *J Geriatr Oncol.* 2015 Jan;6(1):8-14.
6. Naciye Vardar Yagli, Ozlem Ulger. The effects of yoga on the quality of life and depression in elderly breast cancer patients. *Complement Ther Clin Pract.* 2015 Feb;21(1):7-10. doi: 10.1016/j.ctcp.2015.01.002. Epub 2015 Jan 29.
7. Vardar Yağılı N, Şener G, Arıkan H et al. Do yoga and aerobic exercise training have impact on functional capacity, fatigue, peripheral muscle strength, and quality of life in breast cancer survivors? *Integr Cancer Ther.* 2015 Mar;14(2):125-32. doi: 10.1177/1534735414565699. Epub 2015 Jan 6.
8. Rao RM, Raghuram N, Nagendra HR, et al. Effects of an integrated Yoga Program on Self-reported Depression Scores in Breast Cancer Patients Undergoing Conventional Treatment: A Randomized Controlled Trial. *Indian J Palliat Care.* 2015 May-Aug;21(2):174-81. doi: 10.4103/0973-1075.156486.
9. Zetzl T, Renner A, Pitting A et al. Yoga effectively reduces fatigue and symptoms of depression in patients with different types of cancer. *Support Care Cancer* 2020 Oct 7 doi:10.10007/s00520-020-05794-2

Qigong

1. Zeng Y, Xie X, Cheng ASK (2019) Qigong or Tai Chi in Cancer Care: an Updated Systematic Review and Meta-analysis. *Curr Oncol Rep;*21(6):48. doi: 10.1007/s11912-019-0786-2.

2. Fong SS, Choi AW, Luk WS: Bone Mineral Density, Balance Performance, Balance Self-Efficacy, and Falls in Breast Cancer Survivors With and Without Qigong Training. *Integr Cancer Ther.* 2016 Dec 1:1534735416686687. doi: 10.1177/1534735416686687.
3. Larkey LK, Roe DJ, Weihs KL et al. (2015) Randomized controlled trial of Qigong/Tai Chi Easy on cancer-related fatigue in breast cancer survivors. *Annals of behavioral medicine : a publication of the Society of Behavioral Medicine.* 49: 165-176 .
4. Zeng Y, Luo T, Xie H et al. Health benefits of qigong or tai chi for cancer patients: a systematic review and meta-analyses. *Complement Ther Med.* 2014 Feb;22(1):173-86.

Tai Chi

1. Irwin M R, Olmstead R, Carrillo C et al: Tai Chi Chih Compared With Cognitive Behavioral Therapy for the Treatment of Insomnia in Survivors of Breast Cancer: A Randomized, Partially Blinded, Noninferiority Trial . *J Clin Oncol.* 2017 Aug 10;35(23):2656-2665. doi: 10.1200/JCO.2016.71.0285. Epub 2017 May 10.
2. Larkey LK, Roe DJ, Weihs KL et al. (2015) Randomized controlled trial of Qigong/Tai Chi Easy on cancer-related fatigue in breast cancer survivors. *Annals of behavioral medicine : a publication of the Society of Behavioral Medicine.* 49: 165-176 .
3. Pan Y, Yang K, Shi X, et al: Tai chi chuan exercise for patients with breast cancer: a systematic review and meta-analysis. *Evid Based Complement Alternat Med.* 2015;2015:535237.
4. Yan JH, Pan L, Zhang XM et al. Lack of efficacy of Tai Chi in improving quality of life in breast cancer survivors: a systematic review and meta-analysis. *Asian Pac J Cancer Prev.* 2014;15(8):3715-20.

Hypnosis

1. Cramer H, Lauche R, Paul A et al. Hypnosis in Breast Cancer Care: A Systematic Review of Randomized Controlled Trials. *Integr Cancer Ther.* 2015 Jan;14(1):5-15. Epub 2014 Sep 18.
2. Montgomery GH, Schnur JB, Kravits K. Hypnosis for cancer care: Over 200 years young. *CA Cancer J Clin.* 2012 Nov 20. doi: 10.3322/caac.21165.



Komplementäre Therapien

Rezidivprävention / Verbesserung Gesamtüberleben I

Beeinflussbare Lebensstilfaktoren – Sport – Genussmittel

	Oxford		
	LoE	GR	AGO
■ Körperliches Training/Sport (das Äquivalent zu 3–5 Std. mäßiggradigem „Walking“ verbessert DFS und OS und kardiopulmonale Funktion)	2a	A	++
■ Nikotinreduktion	2b	A	+
■ Alkoholkonsum reduzieren (< 6g/die)	2b	A	+

Physical exercise

1. Hong, F., Ye, W., Kuo, C. H., et al.: (2019). Exercise Intervention Improves Clinical Outcomes, but the “Time of Session” is Crucial for Better Quality of Life in Breast Cancer Survivors: A Systematic Review and Meta-Analysis. *Cancers*, 11(5), 706.
2. Nechuta S, Chen WY, Cai H: A pooled analysis of post-diagnosis lifestyle factors in association with late estrogen-receptor-positive breast cancer prognosis. *Int J Cancer*. 2016 May 1;138(9):2088-97. doi: 10.1002/ijc.29940.
3. de Glas NA, Fontein DB, Bastiaannet E, et al: Physical activity and survival of postmenopausal, hormone receptor-positive breast cancer patients: Results of the tamoxifen exemestane adjuvant multicenter lifestyle study. *Cancer* 2014;120:2847-2854.
4. Friedenreich CM, Neilson HK, Woolcott CG, et al: Inflammatory Marker Changes in a Yearlong Randomized Exercise Intervention Trial among Postmenopausal Women. *Cancer Prev Res (Phila)*. 2012 Jan;5(1):98-108.
5. Zeng H, Irwin ML, Lu L, et al: Physical activity and breast cancer survival: an epigenetic link through reduced methylation of a tumor suppressor gene L3MBTL1. *Breast Cancer Res Treat*. 2012 May;133(1):127-35. doi: 10.1007/s10549-011-1716-7. Epub 2011 Aug 12.

Improvements in DFS and OS, prevention of recurrence

1. Patel AV, Friedenreich CM, Moore SC et al. (2019) American College of Sports Medicine Roundtable Report on Physical Activity, Sedentary Behavior, and Cancer Prevention and Control. *Med Sci Sports Exerc*;51(11):2391-2402.
2. Jung, A. Y., Behrens, S., Schmidt, M et al. (2019). Pre-to postdiagnosis leisure-time physical activity and prognosis in

- postmenopausal breast cancer survivors. *Breast Cancer Research*, 21(1), 117.
3. Borch KB, Braaten T, Lund E et al. Physical activity before and after breast cancer diagnosis and survival - the Norwegian women and cancer cohort study. *BMC Cancer*. 2015 Dec 16;15(1):967. doi: 10.1186/s12885-015-1971-9.
 4. Lahart IM, Metsios GS, Nevill AM, et al. Physical activity, risk of death and recurrence in breast cancer survivors: A systematic review and meta-analysis of epidemiological studies. *Acta Oncol*. 2015 May;54(5):635-54.
 5. Zhong S, Jiang T, Ma T et al. Association between physical activity and mortality in breast cancer: a meta-analysis of cohort studies. *Eur J Epidemiol*. 2014 Jun;29(6):391-404.

Smoking

1. Pierce JP, Patterson RE, Senger C et al: Lifetime cigarette smoking and breast cancer prognosis in the after breast cancer pooling project. *J Natl Cancer Inst* 2014;106:djt359.
2. Wang K, Li F, Zhang X: Smoking increases risks of all-cause and breast cancer specific mortality in breast cancer individuals: a dose-response meta-analysis of prospective cohort studies involving 39725 breast cancer cases. *Oncotarget*. 2016 Dec 13;7(50):83134-83147. doi: 10.18632/oncotarget.13366.
3. Pierce JP, Patterson RE, Senger CM: Lifetime cigarette smoking and breast cancer prognosis in the After Breast Cancer Pooling Project. *J Natl Cancer Inst*. 2014 Jan;106(1):djt359. doi: 10.1093/jnci/djt359.
4. Bérubé S, Lemieux J, Moore L: Smoking at time of diagnosis and breast cancer-specific survival: new findings and systematic review with meta-analysis. *Breast Cancer Res*. 2014 Apr 19;16(2):R42. doi: 10.1186/bcr3646.

Alcohol

1. Larsen SB, Kroman N, Ibfelt EH: Influence of metabolic indicators, smoking, alcohol and socioeconomic position on mortality after breast cancer. *Acta Oncol*. 2015 May;54(5):780-8. doi: 10.3109/0284186X.2014.998774.
2. Simapivapan P, Boltong A, Hodge A.: To what extent is alcohol consumption associated with breast cancer recurrence and second primary breast cancer?: A systematic review. *Cancer Treat Rev*. 2016 Nov;50:155-167. doi: 10.1016/j.ctrv.2016.09.010.
3. Katzke VA, Kaaks R, Kühn T. Lifestyle and cancer risk. *Cancer J*. 2015 Mar-Apr;21(2):104-10.
4. Jayasekara H, MacInnis RJ, Room R, English DR. Long-Term Alcohol Consumption and Breast, Upper Aero-Digestive Tract and Colorectal Cancer Risk: A Systematic Review and Meta-Analysis. *Alcohol Alcohol*. 2015 Sep 22. pii: agv110. [Epub ahead of print]
5. Choi YJ, Jyung SK, Lee JH. Ligth Alcohol Drinking and Risko of Cancer: A Meta-Analysis of Cohort Studies. *Cancer Res Treat* 2018;50:474-487.

Komplementäre Therapien

Rezidivprävention / Verbesserung Gesamtüberleben II

Beeinflussbare Lebensstilfaktoren – Ernährung

	Oxford		
	LoE	GR	AGO
■ Anstreben eines normalen BMI	1a	A	++
■ Ernährung mit geringem Fettanteil (Ernährungsberatung empfohlen)	1a	B	+
■ Ballaststoffhaltige Lebensmittel (u.a. Saaten, z.B. Leinsamen)	2a	B	+
■ Beachten genereller Ernährungsempfehlungen (z.B. von DGE, WCRF) im Sinne einer mediterranen (Vollwert-)Ernährung	2a	B	++
■ Diät-Extreme	2a	B	--

Adherence to normal body weight/BMI

1. Mei L, He L, Song Y et al. Association between obesity with disease-free survival and overall survival in triple-negative breast cancer. A meta-analysis. Medicine 2018;97:19
2. (Brenner DR, Brockton NT, Kotsopoulos J: Breast cancer survival among young women: a review of the role of modifiable lifestyle factors. Cancer Causes Control. 2016 Apr;27(4):459-72. doi: 10.1007/s10552-016-0726-5.)
3. (Dieli-Conwright CM, Lee K, Kiwata JL: Reducing the Risk of Breast Cancer Recurrence: an Evaluation of the Effects and Mechanisms of Diet and Exercise. Curr Breast Cancer Rep. 2016;8(3):139-150.)
4. (Schwingshackl L, Hoffmann G: Adherence to Mediterranean diet and risk of cancer: an updated systematic review and meta-analysis of observational studies. Cancer Med. 2015 Dec;4(12):1933-47. doi: 10.1002/cam4.539.)
5. (Ferrini K, Ghelfi F, Mannucci R: Lifestyle, nutrition and breast cancer: facts and presumptions for consideration. Ecancermedicalscience. 2015 Jul 23;9:557. doi: 10.3332/ecancer.2015.557. eCollection 2015.)
6. Rock CL, Flatt SW, Byers TE et al. Results of the Exercise and Nutrition to Enhance Recovery and Good Health for You (ENERGY) Trial: A Behavioral Weight Loss Intervention in Overweight or Obese Breast Cancer Survivors. J Clin Oncol. 2015 Oct 1;33(28):3169-76. doi: 10.1200/JCO.2015.61.1095. Epub 2015 Aug 17.
7. Ligibel JA, Cirrincione CT, Liu M2 et al. Body Mass Index, PAM50 Subtype, and Outcomes in Node-Positive Breast Cancer: CALGB 9741 (Alliance). J Natl Cancer Inst. 2015 Jun 25;107(9). pii: djv179. doi: 10.1093/jnci/djv179. Print 2015 Sep.

8. Jiralerspong S, Goodwin PJ: Obesity and Breast Cancer Prognosis: Evidence, Challenges, and Opportunities. *J Clin Oncol* , 2016; 34:4203-4216
9. Nechuta S, Chen WY, Cal H et al.: A pooled analysis of post-diagnosis lifestyle factors in association with late estrogen-receptor positive breast cancer prognosis. *Int J Cancer*. 2016 May 1; 138(9): 2088–2097. doi:10.1002/ijc.29940
10. Lauby-Secretan, B., et al.: Body Fatness and Cancer--Viewpoint of the IARC Working Group. *N Engl J Med*, 2016. 375(8): p. 794-8.

Übergewicht

1. Mei L, He L, Song Y et al. Association between obesity with disease-free survival and overall survival in triple-negative breast cancer. A meta-analysis. *Medicine* 2018;97:19

Low-Fat Diet

1. De Cicco P, Catani MV et al. Nutrition and Breast Cancer: A Literature Review on Prevention, Treatment and Recurrence. *Nutrients* 2019 Jul 3;11(7).
2. Chlebowski R, Anderson G et al. Low-Fat Dietary Pattern and Cancer Mortality in the Women's Health Initiative (WHI) Randomized Controlled Trial, *JNCI Cancer Spectrum*, Volume 2, Issue 4, October 2018
3. Dieli-Conwright CM, Lee K, Kiwata JL: Reducing the Risk of Breast Cancer Recurrence: an Evaluation of the Effects and Mechanisms of Diet and Exercise. *Curr Breast Cancer Rep.* 2016;8(3):139-150.
4. Chlebowski RT, Aragaki AK, Anderson GL: Low-Fat Dietary Pattern and Breast Cancer Mortality in the Women's Health Initiative Randomized Controlled Trial. *J Clin Oncol.* 2017 Sep 1;35(25):2919-2926. doi: 10.1200/JCO.2016.72.0326. Epub 2017 Jun 27.
5. Makarem N, Chandran U et al. Dietary Fat in Breast Cancer Survival. *Annu Rev Nutr.* 2013 ; 33
6. Xing MY, Xu SZ, Shen P: Effect of low-fat diet on breast cancer survival: A meta-analysis. *Asian Pacific journal of cancer prevention : APJCP* 2014;15:1141-1144.

Fiber

1. McQuade JL et al. Modulating the microbiome to improve therapeutic response in cancer *Lancet Oncol.* 2019; 20: e77-e91
2. De Cicco P, Catani MV et al. Nutrition and Breast Cancer: A Literature Review on Prevention, Treatment and Recurrence. *Nutrients* 2019 Jul 3;11(7).
3. World Cancer Research Fund International/American Institute for Cancer Research Continuous Update Project Report: Diet,

Nutrition, Physical Activity, and Breast Cancer Survivors. 2014. Available at: www.wcrf.org/sites/default/files/Breast-Cancer-Survivors-2014-Report.pdf

Adherence to general nutrition – guidelines:

1. De Cicco P, Catani MV et al. Nutrition and Breast Cancer: A Literature Review on Prevention, Treatment and Recurrence. *Nutrients* 2019 Jul 3;11(7).
2. Hou R, Wei J et al. Healthy dietary pattern and risk and survival of breast cancer: a meta-analysis of cohort studies, *Cancer causes & control* (2019) 30: 835-846
3. World Cancer Research FundInternational/American Institute for Cancer Research, Continous Update Project: Diet, Nutrition, Physical Activity, and Breast Cancer Survivors. 2014. Available at: www.wcrf.org/sites/default/files/Breast-Cancer-Survivors-2014-Report.pdf
4. Ferrini K, Ghelfi F, Mannucci R: Lifestyle, nutrition and breast cancer: facts and presumptions for consideration. *Ecancermedicalscience*. 2015 Jul 23;9:557. doi: 10.3332/ecancer.2015.557. eCollection 2015.
5. Gonzales, J. F., N. D. Barnard, D. J. Jenkins, A. J. Lanou, B. Davis, G. Saxe and S. Levin (2014). "Applying the precautionary principle to nutrition and cancer." *J Am Coll Nutr* 33(3): 239-246.
6. Schwingshackl L, Schwedhelm C: Adherence to Mediterranean Diet and Risk of Cancer: An Updated Systematic Review and Meta-Analysis. *Nutrients* 2017 (9) 1063; doi:10.3390/nu9101063
7. Toledo, E., et al., Mediterranean Diet and Invasive Breast Cancer Risk Among Women at High Cardiovascular Risk in the PREDIMED Trial: A Randomized Clinical Trial. *JAMA Intern Med*, 2015. 175(11): p. 1752-60
8. Biasini C, di Nunzio C, Cordani MR, et al.: Mediterranean Diet influences breast cancer relapse: preliminary results of the SETA PROJECT. *Journal of Clinical Oncology*, 2016 ASCO Annual Meeting (June 3-7, 2016).Vol 34, No 15_suppl (May 20 Supplement), 2016: e13039
9. Amireault, S, Fon AJ, Sabiston CM. Promoting Healthy Eating and Physical Activity Behaviors: a systematic review of multiple health behavior change interventions among cancer survivors. *American Journal of lifestyle Medicine* 2016;12:184-199.
- 10.Rhee J, Mattei J, Huges M et al. Diabetes risk reduction diet score. SABSC 2020

Diät Extreme:

1. Mohsen M, Katsiki N et al. Lower carbohydrate diets and all-cause and cause-specific mortality: a population-based cohort study and pooling of prospective studies. *European Heart Journal* 2019, 40, 2870-2879

2. Huebner J., Marienfeld S. et al.: Counseling Patients on Cancer Diets: A Review of the Literature and Recommendations for Clinical Practice. *Anticancer Res.* 2014 Jan; 34(1):39-48.
3. Erickson, N., Boscheri, A., Linke, B. et al.: Systematic review: isocaloric ketogenic dietary regimes for cancer patients. *Med Oncol* (2017) 34: 72. <https://doi.org/10.1007/s12032-017-0930-5>
4. Jochems SHJ, van Osch FH, Bryan RT, et al., Impact of dietary patterns and the main food groups on mortality and recurrence in cancer survivors: a systematic review of current epidemiological literature *BMJ Open* 2017;8:e014530.



Komplementäre Therapien

Rezidivprävention / Verbesserung Gesamtüberleben III

Pflanzliche Therapieansätze – Nahrungsergänzung

	Oxford	GR	AGO
	LoE		
▪ Nach Systemtherapie – Vitamine/Antioxidanzien scheinen nicht mit einem erhöhtem Rezidivrisiko assoziiert	2b	B	+/-
▪ Raucher haben ein höheres Bronchial-Ca-Risiko unter Antioxidanzien	1b	A	-
Prävention eines brustkrebsassoz. Rezidis			
▪ Antioxidanzien	2a	B	+/-
▪ Vitamine (zusätzlich zu ausgewogener Ernährung; Vit. C, E, D)	2a	B	+/-
▪ Sojaprodukte (Phytoöstrogene)	2a	B	+/-
▪ Phytoöstrogene Konzentration ≥ 100 mg Isoflavone pro Tag	2a	B	-
▪ Traubensilberkerze (<i>Cimicifuga racemosa</i>)	3b	C	+/-
▪ Antioxidative Supplamente nach Beendigung der Radiotherapie	2b	B	+/-
▪ Grüner Tee	3a	C	+/-
▪ Orthomolekulare Substanzen (Selen, Zink ...)	5	D	-
▪ Karotenoide erscheinen mit schlechterem Ergebnis assoziiert	2b	B	-
▪ Proteolytische Enzyme (Papain, Trypsin, Chymotrypsin)	3b	B	-
▪ Mistellektine (<i>Viscum album</i>)	1b	C	-
▪ Thymuspeptide (Einfluss auf Überleben)	2a	B	-
▪ Sauerstoff- und Ozon-Therapie	5	D	--
▪ Laetrire (Aprikosenkernerextrakt)	1c	D	--
▪ Methadon	5	D	--
▪ Cancer bush (<i>Sutherlandia frutescens</i>), Devil's claw (<i>Harpagophytum procumbens</i>), Rooibos Tee (<i>Aspalathus linearis</i>), Bambara-Erdnuss (<i>Vignea subterranean</i>)	5	D	-
▪ Weihrauch	5	D	-

General

1. Hervik JB, Stub T: Adverse effects of non-hormonal pharmacological interventions in breast cancer survivors, suffering from hot flashes: A systematic review and meta-analysis. *Breast Cancer Res Treat.* 2016 Nov;160(2):223-236.

Post treatment vitamin and/or antioxidant supplements

1. Drewe J, Bucher KA, Zahner C. A systematic review of non-hormonal treatments of vasomotor symptoms in climacteric and cancer patients. *Springerplus.* 2015 Feb 10;4:65. doi: 10.1186/s40064-015-0808-y. eCollection 2015.
2. Sodde VK, Lobo R, Kumar N, et al. Cytotoxic activity of Macrosolen parasiticus (L.) Danser on the growth of breast cancer cell line (MCF-7). *Pharmacogn Mag.* 2015 May;11(Suppl 1):S156-60. doi: 10.4103/0973-1296.157719.
3. Salarabadi A, Bidgoli SA, Madani SH. Roles of Kermanshahi Oil, Animal Fat, Dietary and Non- Dietary Vitamin D and other Nutrients in Increased Risk of Premenopausal Breast Cancer: A Case Control Study in Kermanshah, Iran. *Asian Pac J Cancer Prev.* 2015;16(17):7473-8.
4. Karimi Z, Bahadoran Z, Abedini S, et al. Dietary total antioxidant capacity and the risk of breast cancer: a case-control study. *East Mediterr Health J.* 2015 Sep 28;21(8):564-71.
5. Carioca AA, Verde SM, Luzia LA, et al. Association of oxidative stress biomarkers with adiposity and clinical staging in women with breast cancer. *Eur J Clin Nutr.* 2015 Nov;69(11):1256-61. doi: 10.1038/ejcn.2015.84. Epub 2015 Jun 3.

6. Martin-Herranz A, Salinas-Hernández P. Vitamin D supplementation review and recommendations for women diagnosed with breast or ovary cancer in the context of bone health and cancer prognosis/risk. *Crit Rev Oncol Hematol.* 2015 Oct;96(1):91-9. doi: 10.1016/j.critrevonc.2015.05.006. Epub 2015 May 19.

Soy as normal part of the diet/soy concentrates

1. Wu AH, Spicer D, Garcia A, et al. Double-Blind Randomized 12-Month Soy Intervention Had No Effects on Breast MRI Fibroglandular Tissue Density or Mammographic Density. *Cancer Prev Res (Phila).* 2015 Oct;8(10):942-51. doi: 10.1158/1940-6207.CAPR-15-0125. Epub 2015 Aug 14.
2. Fritz H, Seely D, Flower G, et al.: Soy, red clover, and isoflavones and breast cancer: A systematic review. *PLoS one* 2013;8:e81968.

Black cohosh

1. Fritz H, Seely D, McGowan J, et al: Black cohosh and breast cancer: A systematic review. *Integrative cancer therapies* 2014;13:12-29.

Laetrile treatment for cancer

1. Milazzo S, Ernst E, Lejeune S, et al.: Laetrile treatment for cancer. *Cochrane Database of Systematic Reviews* 2011, Issue 11. Art. No.: CD005476. DOI: 10.1002/14651858.CD005476.pub3.

St John's Wort

1. Caraci F, Crupi R, Drago F, et al. Metabolic drug interactions between antidepressants and anticancer drugs: focus on selective serotonin reuptake inhibitors and hypericum extract. *Curr Drug Metab.* 2011 Jul 1;12(6):570-7.
2. Schellander R, Donnerer J: Antidepressants: clinically relevant drug interactions to be considered. *Pharmacology.* 2010;86(4):203-15. Epub 2010 Sep 8.
3. Nahrstedt A, Butterweck V: Lessons learned from herbal medicinal products: the example of St. John's Wort (perpendicular). *J Nat Prod.* 2010 May 28;73(5):1015-21.

Red clover

1. Fritz H, Seely D, Flower G et al. Soy, red clover, and isoflavones and breast cancer: A systematic review. *PLoS One.* 2013 Nov 28;8(11):e81968.

2. Geller SE, Shulman LP, van Breemen RB et al.: Safety and efficacy of black cohosh and red clover for the management of vasomotor symptoms: a randomized controlled trial. *Menopause*. 2009;16(6):1156–1166.

Dong Quai

1. Zhuang SR, Chiu HF, Chen SL, Effects of a Chinese medical herbs complex on cellular immunity and toxicity-related conditions of breast cancer patients. *Br J Nutr.* 2011 Aug 25:1-7.
2. Rotem C, Kaplan B: Phyto-Female Complex for the relief of hot flushes, night sweats and quality of sleep: randomized, controlled, double-blind pilot study. *Gynecol Endocrinol.* 2007;23(2):117-122.

Mistletoe

1. Freudig M., Keinki C., Micke O., Mistletoe in oncological treatment: a systematic review. *Journal of Cancer Research and Clinical Oncology* (2019) 145:695–707

Ginseng root

1. Yamada N, Araki H, Yoshimura H: Identification of antidepressant-like ingredients in ginseng root (*Panax ginseng* C.A. Meyer) using a menopausal depressive-like state in female mice: participation of 5-HT2A receptors. *Psychopharmacology (Berl)*. 2011 Aug;216(4):589-99.
2. Peralta EA, Murphy LL, Minnis J, et al.: American Ginseng inhibits induced COX-2 and NFkB activation in breast cancer cells. *J Surg Res.* 2009 Dec;157(2):261-7.

Bromelain+Papain+Selen+Lektin bei AI-induced athralgia

1. Uhlenbruck B, Van Leendert R, Schneider B et al.: Reduced side-effects of adjuvant hormone therapy in breast cancer patients by complementary medicine. *In Vivo.* 2010 Sep-Oct;24(5):799-802.

Green Tea

1. Gianfredi V, Nucci D, Abalsamo A, et al. Green Tea consumption and risk of breast cancer and recurrence – a systematic review and meta-analysis of observational studies. *Nutrients* 2018;10;pii:E1886.
2. Najaf Najafi M, Salehi M. et al. The association between green tea consumption and breast cancer risk. A systematic review and meta-analysis. *Phytother Res* 2018;32:1855-1864.