



Diagnostik und Therapie früher und fortgeschritten Mammakarzinome

Brustkrebs Nachsorge



Brustkrebs Nachsorge

- **Versionen 2002–2020:**

Bauerfeind / Bischoff / Blöhmaier / Böhme / Costa / Diel / Friedrich /
Gerber / Hanf / Heinrich / Huober / Janni / Kaufmann / Kolberg-
Liedtke / Kümmel / Lux / Maass / Möbus / Müller-Schimpfle/
Mundhenke / Oberhoff / Rody / Scharl / Solbach / Solomayer /
Thomssen / Wöckel

- **Version 2021:**

Gluz / Lüftner

Aktualisierung der Therapieempfehlungen nach Durchsicht der ASCO und ACS Guidelines*, sowie der S3 Leitlinie
Durchgeführte „Medline“ Suche nach „Surveillance Breast Cancer“ und „Follow up primary breast cancer“ (2018/01-2021/01)

*Runowicz CD et al., American Cancer Society/American Society of Clinical Oncology Breast Cancer Survivorship Care Guideline, JCO 34 :611-635,



Brustkrebs Nachsorge

Ziele

	Oxford		
	LoE	GR	AGO
Früherkennung von heilbaren Rezidiven			
▪ Intramammäre Rezidive	1a	B	++
▪ Lokoregionäre Rezidive*	1a	B	++
Früherkennung kontralateraler Karzinome	1a	B	++
Früherkennung von Metastasen			
▪ Früherkennung symptomatischer Metastasen	3b	C	+
▪ Früherkennung asymptomatischer Metastasen	1a	A	-

* Das lokoregionäre Rezidiv ist mit einem erhöhten Mortalitätsrisiko bei nodalpositiven, PR-negativen, jüngeren Patientinnen und einem kurzen Zeitintervall von Erstdiagnose bis Rezidiv verbunden.

1. De Bock GH, Bonnema J, van Der Hage J et al., Effectiveness of Routine Visits and Routine Tests in Detecting Isolated Locoregional Recurrences After Treatment for Early-Stage Invasive Breast Cancer: A Meta-Analysis and Systematic Review. *J Clin Oncol* 2004; 22 (19): 4010-4018.
2. Margenthaler JA, Allan E, Cheng L, et al.. Surveillance of Patients With Breast Cancer After Curative-Intent Primary Treatment: Current Practice Patterns. *Journal of Oncology Practice* 2012; 8(2): 79 – 83.
3. Parmeshwar R, Margenthaler JA, Allam E, et al.. Patient Surveillance After Initial Breast cancer Therapy Variation by Physician Specialty. *Am J Surg* 2013; 206(2): 218-222.
4. Jochelson M, Hayes DF, Ganz PA. Surveillance and Monitoring in Breast Cancer Survivors: Maximizing Benefit and Minimizing Harm. *ASCO Educational Book* 2013 e13 – e18.
5. Khatcheressian JL, Hurley P, Bantug E, et al.. Breast Cancer Follow-up and Management After Primary Treatment: American Society of Clinical Oncology Clinical Practice Guideline Update . *J Clin Oncol*. 2013 March 1; 31(7):961-965.
6. Runowicz CD, Leach CR, Henry NL, et al. American Cancer Society/American Society of Clinical Oncology Breast Cancer Survivorship Care Guideline. *Journal of Clinical Oncology* 2016;34:611-35.
7. Cardoso F, Kyriakides S, Ohno S, et al. Early breast cancer: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Annals of Oncology* 2019;30:1194-220.
8. Lee et al. *Cancer Res Treat*. 2015 Oct;47(4):765-73. Reduced mortality according to regular follow up **Impact on Survival of Regular**

Postoperative Surveillance for Patients with Early Breast Cancer

9. Moschetti I, Cinquini M, Lambertini M et al., Follow-up strategies for women treated for early breast cancer. Cochrane Database Syst Rev. 2016 May 27;(5):1CD001768.
10. NCCN Clinical Practice Guidelines in Oncology, Breast Cancer Version 3.17-10.17;
https://www.nccn.org/professionals/physician_gls/pdf/breast.pdf
11. NCCN Clinical Practice Guidelines in Oncology, Breast Cancer Version 6.2020
https://www.nccn.org/professionals/physician_gls/pdf/breast_blocks.pdf

Statement: risk factors of mortality after loco-regional recurrence

1. Dent R, Valentini H, Hanna W, et al.. Factors associated with breast cancer mortality after local recurrence. Curr Oncol 2014; 21 (3): e418-25.



Brustkrebs Nachsorge

Ziele

Oxford		
LoE	GR	AGO
2b	B	+
2a	B	+
2b	B	+
3b	B	+

Verbesserung der Lebensqualität

Verbesserung der körperlichen Leistungsfähigkeit

Reduktion bzw. zeitnahe Erkennung therapiebedingter Nebenwirkungen (wie z.B. Osteoporose, Herzinsuffizienz, Fatigue, Neuro-toxizität, Lymphödeme, sexuelle Beschwerden, kognitive Einschränkungen, Sterilität, Zweitmalignome) und Veranlassung notwendiger Therapien

Teilnahme an Interventionsprogrammen in der Nachsorge (z.B. Lifestyle, Therapieadhärenz etc.)

Statement: Outcome measurements

1. Ong WL, Schouwenburg MG, van Bommel ACM et al.. A Standard Set of Value-Based Patient-Centered Outcomes for Breast Cancer: The International Consortium for Health Outcomes Measurement (ICHOM) Initiative. *JAMA Oncol.* 2017 May 1;3(5):677-685.
2. Browall M, Forsberg C, Wengström Y. Assessing patient outcomes and cost-effectiveness of nurse-led follow-up for women with breast cancer - have relevant and sensitive evaluation measures been used? *J Clin Nurs.* 2017 Jul;26(13-14):1770-1786.
3. Cheng KKF, Lim YTE, Koh ZM et al. Home-based multidimensional survivorship programmes for breast cancer survivors. *Cochrane Database Syst Rev.* 2017 Aug 24;8:CD011152.

Statement: Obesity, physical activity and quality of life

1. Bicego D, Brown K. Effects of Exercise on Quality of Life in Women Living with Breast Cancer: A Systematic Review. *The Breast Journal* 2009; 15(1): 45-51.
2. Carson JW, Carson KM, Porter LS et al.. Yoga of Awareness program for menopausal symptoms in breast cancer survivors: results from a randomized trial. *Support Care Cancer* 2009; 17: 1301-1309.
3. Vaskuil DW, van Nes JG, Jungeburgt JM et al.. Maintenance of physical activity and body weight in relation to subsequent quality of life in postmenopausal breast cancer patients. *Annals of Oncology* 2010; 21: 2094–2101.
4. Cramp F, Byron-Daniel J. Exercise for the management of cancer-related fatigue in adults. *Cochrane Database Syst Rev.* 2012 Nov 14;11:CD006145.
5. Bower JE, Garet D, Sternlieb B et al.. Yoga for persistent fatigue in breast cancer survivors: A randomized controlled trial. *Cancer*

- 2012; 118(15): 3766-75.
6. Cramer H, Lange S, Klose P et al.. Can yoga improve fatigue in breast cancer patients? A systematic review. *Acta Oncol* 2012; 51(4): 559 – 60.
 7. Yang H, Brand JS, Fang F et al.. Time-dependent risk of depression, anxiety, and stress-related disorders in patients with invasive and in situ breast cancer. *Int J Cancer*. 2017 Feb 15;140(4):841-852.
 8. Nagy AC, GulAcsi-BArdos P, CserEp Z et al..Late cardiac effect of anthracycline therapy in physically active breast cancer survivors - a prospective study. *Neoplasma*. 2017;64(1):92-100.
 9. Baumann FT, Bieck O, Oberste M et al.. Sustainable impact of an individualized exercise program on physical activity level and fatigue syndrome on breast cancer patients in two German rehabilitation centers. *Support Care Cancer*. 2017 Apr;25(4):1047-1054.
 10. NCCN-Guidelines Version 1.2017. Cancer-related Fatigue.
 11. Fabi A, Bhargava R, Fatigoni S, et al. Cancer-related fatigue: ESMO Clinical Practice Guidelines for diagnosis and treatment. *Annals of Oncology* 2020;31:713-23.
 12. Curigliano G, Lenihan D, Fradley M, et al. Management of cardiac disease in cancer patients throughout oncological treatment: ESMO consensus recommendations. *Annals of Oncology* 2020;31:171-90.

Statement: Obesity and breast cancer prognosis

1. Ewertz M, Jensen MB, Gunnarsdóttir KÁ et al.. Effect of obesity on prognosis after early-stage breast cancer. *J Clin Oncol* 2011; 29(1): 25-31.
2. Cespedes Feliciano EM, Kroenke CH, Bradshaw PT et al..Postdiagnosis Weight Change and Survival Following a Diagnosis of Early-Stage Breast Cancer. *Cancer Epidemiol Biomarkers Prev*. 2017 Jan;26(1):44-50.

Statement: Lymphedema

1. Soran A, Ozmen T, McGuire KP et al..The importance of detection of subclinical lymphedema for the prevention of breast cancer-related clinical lymphedema after axillary lymph node dissection; a prospective observational study.*Lymphat Res Biol* 2014;12(4): 289-9.
2. Basta MN, Wu LC, Kanchwala SK et al.. Reliable prediction of postmastectomy lymphedema: the Risk Assessment Tool Evaluating Lymphedema. *Am J Surg*. 2017 Jun;213(6):1125-1133.
3. Ferguson CM, Swaroop MN, Horick N, et al. Impact of Ipsilateral Blood Draws, Injections, Blood Pressure Measurements, and Air Travel on the Risk of Lymphedema for Patients Treated for Breast Cancer. *Journal of Clinical Oncology* 2016;34:691-8.

Statement: sexual disorders and cognitive impairment:

1. Runowicz CD, Leach CR, Henry L et al.. American Cancer Society/American Society of Clinical Oncology breast cancer survivorship care guideline. CA Cancer J Clin 2016; 66: 43-73.
2. Janelsins MC, Heckler CE, Peppone LJ et al.. Cognitive Complaints in Survivors of Breast Cancer After Chemotherapy Compared With Age-Matched Controls: An Analysis From a Nationwide, Multicenter, Prospective Longitudinal Study. J Clin Oncol. 2017 Feb 10;35(5):506-514.
3. Bernstein LJ, McCreath GA, Komeylian Z et al..Cognitive impairment in breast cancer survivors treated with chemotherapy depends on control group type and cognitive domains assessed: A multilevel meta-analysis. Neurosci Biobehav Rev. 2017 Dec;83:417-428.

Statement: Secondary tumors:

1. Hoekstra N, Fleury E, Merino Lara TR, et al. Long-term risks of secondary cancer for various whole and partial breast irradiation techniques. Radiother Oncol. 2018 Sep;128(3):428-433



Monitoring nach kardiotoxischer Therapie (z.B. Anthrazyklin; Anti-HER2)

- **Echokardiografie 6-12 Monate und nach 5 Jahren nach der Beendigung der Therapie** (v.a. Im Fall der höher dosierten Anthrazykline oder im Fall der vorliegenden Risikofaktoren (linksseitige Radiatio, Nikotin, Hypertonie, Diabetes, Dyslipidämie, Übergewicht, Alter ≥ 60 J. während der Therapie, kardiale Vorerkrankung: reduzierte Pumpfunktion, Zustand nach Myokardinfarkt, \geq mittelgradiges Vitium))
- ggf. BNP Messung

1. Curigliano G, Lenihan D, Fradley M, et al. Management of cardiac disease in cancer patients throughout oncological treatment: ESMO consensus recommendations. *Annals of Oncology* 2020;31:171-90.
2. Michel L, Rassaf T, Totzeck M. Biomarkers for the detection of apparent and subclinical cancer therapy-related cardiotoxicity. *J Thorac Dis* 2018;10:S4282-S95.
3. Rassaf T, Totzeck M, Backs J, et al. Onco-Cardiology: Consensus Paper of the German Cardiac Society, the German Society for Pediatric Cardiology and Congenital Heart Defects and the German Society for Hematology and Medical Oncology. *Clin Res Cardiol* 2020;109:1197-222.



Brustkrebs Nachsorge Inhalte und Ziele

	Oxford		
	LoE	GR	AGO
▪ Evaluation laufender adjuvanter Therapien	2b	B	++
▪ inkl. Überprüfung der Adhärenz endokriner Therapien ▪ Überprüfung des Menopausenstatus z.B. bei CT-induzierter Amenorrhoe 6-monatlich (FSH und/oder Blutungsanamnese bei Patientinnen <45 Jahre alt) und ggf. Hinzunahme der ovarien Suppression (bis zu 2 Jahre nach der CT) oder Umstellung der AHT	5	D	++
▪ Pro-aktive Verbesserung der Adhärenz anstreben durch:			
▪ Patientenaufklärung über die günstigen Daten einer 5- bis 10-jährigen adj. endokrinen Therapie ▪ Frühzeitige Therapie von Nebenwirkungen (z.B. Sport-intervention, NSAID, Vitamin D / Calcium-Substitution)			

Evaluation of current adjuvant therapy

1. Hershman DL, Kushi LH, Shao T et al.. Early Discontinuation and Nonadherence to Adjuvant Hormonal Therapy in a Cohort of 8,769 Early-Stage Breast Cancer Patients. *J Clin Oncol* 2010; 28: 4120-4128.
2. Lueck H-J, Hadji P, Harbeck N et al.. 24 Months Follow-Up Results from PACT (Patient's Anastrozole Compliance to Therapy Programme), a Non-Interventional Study Evaluating the Influence of a Standardized Information Service on Compliance in Postmenopausal Women with Early Breast Cancer. *SABCS 2011* [P5-17-05].
3. Neven P, Markopoulos C, Tanner M et al.. The impact of educational materials on compliance and persistence rates with adjuvant aromatase inhibitor treatment: first-year results from the compliance of aromatase inhibitors assessment in daily practice through educational approach (CARIATIDE) study. *Breast*. 2014 Aug;23(4):393-9.
4. Hershman DL, Kushi LH, Hillyer GC et al.. Psychosocial factors related to non persistence with adjuvant endocrine therapy among women with breast cancer: the Breast Cancer Quality of Care Study (BQUAL). *Breast Cancer Res Treat*. 2016 May;157(1):133-43.
5. Goss PE, Ingle JN, Pritchard KI et al.. Extending Aromatase-Inhibitor Adjuvant Therapy to 10 Years. *N Engl J Med*. 2016 Jul 21;375(3):209-19.
6. Nabieva N, Kellner S, Fehm T et al.. Patient and tumor characteristics and their influence on early therapy persistence with letrozole in postmenopausal patients with early breast cancer. *Ann Oncol*. 2017 Oct 10. doi: 10.1093/annonc/mdx630.
7. Laroche F, Perrot S, Medkour T et al.. Quality of life and impact of pain in women treated with aromatase inhibitors for breast cancer. A multicenter cohort study. *PLoS One*. 2017 Nov 8;12(11):e0187165.
8. Kim, H. A., Lee, J. W., Nam, S. J., et al Adding Ovarian Suppression to Tamoxifen for Premenopausal Breast Cancer: A Randomized Phase

III Trial J Clin Oncol 2020 Feb 10;38(5):434-443



Brustkrebs Nachsorge Inhalte und Ziele

	Oxford		
	LoE	GR	AGO
• Psychosoziale Aspekte der Beratung	4	C	+
- Schwangerschaft, Kontrazeption, Sexualität, Lebensqualität, Menopausensyndrom, Angst vor Rezidiv			
- Einbeziehung von Bezugspersonen (Partner, Familie, Freunde, Betreuer)			
• Zweitmeinung zur Primärtherapie	2c	B	++
• Allgemeine Beratung (z.B. Überprüfung der aktuellen Indikation zur genetischen Beratung (z.B. bei neu aufgetretene Mamma-/Ovarial-/Pankreas-/Prostata-Ca. in der Familie), HRT, prophylaktische Operationen, Brustrekonstruktion)	2c	C	+

Statement: Psycho-social aspects

1. Drolet M, Maunsell E, Brisson J et al. Not Working 3 Years After Breast Cancer: Predictors in a Population-Based Study. *J Clin Oncol* 2005; 23(33): 8305-8312.
2. Scheier MF, Helgeson VS, Schulz R et al.. Interventions to Enhance Physical and Psychological Functioning Among Younger Women Who Are Ending Nonhormonal Adjuvant Treatment for Early-Stage Breast Cancer. *J Clin Oncol* 2005; 23(19): 4298-4311.
3. Fors EA, Bertheussen GF, Thune I et al.: Psychosocial interventions as part of breast cancer rehabilitation programs? Results from a systematic review. *Psycho-Oncology* 2011; 20: 909-918.
4. Silva C, Caramelo O, Almeida-Santos T et al.. Factors associated with ovarian function recovery after chemotherapy for breast cancer: a systematic review and meta-analysis. *Hum Reprod*. 2016 Dec;31(12):2737-2749.
5. Luke B, Brown MB, Missmer SA et al.. Assisted reproductive technology use and outcomes among women with a history of cancer. *Hum Reprod*. 2016 Jan;31(1):183-9.
6. Gudenkauf LM, Ehlers SL. Psychosocial interventions in breast cancer survivorship care. *Breast*. 2017 Nov 20;38:1-6.
7. Rogers LQ, Courneya KS, Anton PM et al.. Effects of a multicomponent physical activity behavior change intervention on fatigue, anxiety, and depressive symptomatology in breast cancer survivors: randomized trial. *Psychooncology*. 2017 Nov;26(11):1901-1906.
8. Y Kim, DA Kashy, RL Spillers, et al: Needs assessment of family caregivers of cancer survivors: Three cohorts comparison *Psychooncology* 19:573–582, 2010 Crossref, Medline, Google Scholar
9. Y Kim, RL Spillers, DL Hall: Quality of life of family caregivers 5 years after a relative's cancer diagnosis: Follow-up of the national quality of life survey for caregivers *Psychooncology* 21:273–281, 2012 Crossref, Medline, Google Scholar

10. BA Given, CW Given, PR Sherwood: Family and caregiver needs over the course of the cancer trajectory J Support Oncol 10:57–64,2012 Crossref, Medline, Google Scholar

Statement: prophylactic surgery

1. Rhiem K, Engel C, Graeser M et al.. The risk of contralateral breast cancer in patients from BRCA ½ negative high risk families as compared to patients from BRCA1 or BRCA2 positive families: a retrospective cohort study. Breast Cancer Res. 2012; 14(6): R156.



Brustkrebs Nachsorge

Empfohlene Interventionen

Interventionen hinsichtlich Begleiterkrankungen und Lebensstil, um einen negativen Einfluss auf den Krankheitsverlauf zu reduzieren

	Oxford		
	LoE	GR	AGO
• Einstellung Diabetes mellitus (Typ II) (> 25% unerkannter DM bei postmenopausalem MaCa)	2a	B	++
• Gewichts/Lifestyleintervention (bei BMI <18,5 und >30)	2a	B	+
• Nächtliche Nahrungskaresz > 13h	2b	B	+
• fettreduzierte Diät (mindestens 15 % Kalorienreduktion durch Fett) verbessertes Gesamtnüberleben bei HR-MaCa.)	2b	B	+
• Intervention bei Nikotinabusus (durch Rauchen 2 x erhöhte Brustkrebspezifische, 4 x erhöhte nicht-Brustkrebspezifische Mortalität)	2b	B	++
• Alkoholkonsum reduzieren unter 6 g/d	2b	B	+
• Moderate Sportintervention bei Bewegungsmangel (mind. 150 min/Woche, 2x/Woche)	1b	A	++
• Distress-Reduktion	3b	B	+

1. Onitilo AA, Donald M, Stankowski RV et al.. Breast and prostate cancer survivors in a diabetic cohort: results from the Living with DiabetesStudy. Clin Med Res. 2013 Dec;11(4):210-8.
2. Anderson C, Sandler DP, Weinberg CR et al.. Age- and treatment-related associations with health behavior change among breast cancer survivors. Breast. 2017 Jun;33:1-7.
3. Syrowatka A, Motulsky A, Kurteva S et al.. Predictors of distress in female breast cancer survivors: a systematic review. Breast Cancer Res Treat. 2017 Sep;165(2):229-245. Review.
4. Gudenkauf LM, Ehlers SL. Psychosocial interventions in breast cancer survivorship care. Breast. 2017 Nov 20;38:1-6. Review.
5. Mehra K, Berkowitz A, Sanft T.D et al.. Physical Activity, and Body Weight in Cancer Survivorship. Med Clin North Am. 2017 Nov;101(6):1151-1165. Review
6. Haykowsky MJ, Scott JM, Hudson K et al.. Lifestyle Interventions to Improve Cardiorespiratory Fitness and Reduce Breast Cancer Recurrence. Am Soc Clin Oncol Educ Book. 2017;37:57-64.
7. Chlebowski RT, Aragaki AK, Anderson GL et al. Low-Fat Dietary Pattern and Breast Cancer Mortality in the Women's Health Initiative Randomized Controlled Trial. J Clin Oncol. 2017 Sep 1;35(25):2919-2926.
8. Marinac CR, Nelson SH, Breen CI et al..Prolonged Nightly Fasting and Breast Cancer Prognosis. JAMA Oncol. 2016 Aug 1;2(8):1049-55.
9. Sonnenblick A, Agbor-Tarh D, Bradbury I, et al. Impact of Diabetes, Insulin, and Metformin Use on the Outcome of Patients With Human Epidermal Growth Factor Receptor 2-Positive Primary Breast Cancer: Analysis From the ALTTO Phase III Randomized Trial. Journal of Clinical Oncology 2017;35:1421-9.

Statement: for all statements see most recent literature see at Survivorship care guidelines of ASC and ASCO

1. Runowcz CD, Leach CR, Henry L et al.. American Cancer Society/American Society of Clinical Oncology breast cancer survivorship care guideline. CA Cancer J Clin 2016; 66: 43-73.
2. Rock CL, Doyle C, Demark-Wahnefried W, et al. Nutrition and physical activity guidelines for cancer survivors. CA: a cancer journal for clinicians 2012;62:243-74.

Weight intervention.

1. Chajès V, Romieu I. Nutrition and breast cancer. Maturitas, 2014; 77 (1): 7–11.
2. Shaikh H, Bradhurst P, Ma LX et al.: Body weight management in overweight and obese breast cancer survivors. The Cochrane database of systematic reviews 2020;12:Cd012110.
3. Goodwin PJ, Segal RJ, Vallis M, et al. The LISA randomized trial of a weight loss intervention in postmenopausal breast cancer. npj Breast Cancer 2020;6:6.
4. Janni W, Rack B, Friedl T, et al. Abstract GS5-03: Lifestyle Intervention and Effect on Disease-free Survival in Early Breast Cancer Pts: Interim Analysis from the Randomized SUCCESS C Study. Cancer Research 2019;79:GS5-03-GS5-.

Moderate sport intervention when physical activity was reduced

1. Chlebowski RT. Nutrition and physical activity influence on breast cancer incidence and outcome. Breast 2013; Aug;22 Suppl 2: S30-7.
2. Patsou ED, Alexias GD, Anagnostopoulos FG et al.. Effects of physical activity on depressive symptoms during breast cancer survivorship: a meta-analysis of randomised control trials. ESMO Open. 2017 Dec 11;2(5):e000271
3. Friedenreich CM, Stone CR, Cheung WY, Hayes SC. Physical Activity and Mortality in Cancer Survivors: A Systematic Review and Meta-Analysis. JNCI cancer spectrum 2020;4:pkz080.



AGO e. V.
in der DGGO e.V.
in der DKD e.V.
Guidelines Breast
Version 2021.10

www.ago-online.de
FÖRDERER DER
ÄRZTEBÜCHER
VERLEGEN

Nightly fasting

Prolonged nightly fasting improves prognosis in breast cancer patients

retrospective cohort study:

2413 BC-pat. (no diabetes), nightly fasting more or less than 13 hrs

Fasting < 13 hrs: HR 1.36, 36% increase of risk for recurrence
 HR 1.21, n.s. increase of risk for mortality

every 2-hrs-prolonged fasting was correlated with a 20% increase
of sleeping duration

Marinac CR, Nelson SH, Breen CJ et al. JAMA Oncol 2016; 2:1049-1055



Routine-Nachsorgeuntersuchungen bei asymptomatischen Patientinnen

	Oxford		
	LoE	GR	AGO
Untersuchungen			
▪ Anamnese (spezifische Symptome)	1a	A	++
▪ Untersuchung	1a	B	++
▪ Brust-Selbst-Untersuchung	5	D	+
▪ Mammographie	1a	A	++
▪ Mammasonographie	2a	B	++
▪ Mamma-MR in der Routine*	3a	B	+/-
▪ Mamma-MR bei unklarer Mammographie/-sonographie	3b	B	+
▪ Gynäkologische Untersuchung	5	D	++
▪ DXA-Scan zu Therapiebeginn und risikoadaptiert in regelmäßigen Abständen bei Frauen mit frühzeitiger Menopause und Frauen unter AI-Therapie	5	D	+

* Bei erhöhtem Risiko erwägen (Alter unter 50 J., HR+, Beurteilbarkeit in MG+US C/D)

1. Margenthaler JA, Allan E, Cheng L, et al.. Surveillance of Patients With Breast Cancer After Curative-Intent Primary Treatment: Current Practice Patterns. *Journal of Oncology Practice* 2012; 8(2): 79 – 83.
2. Parmeshwar R, Margenthaler JA, Allam E et al.. Patient Surveillance After Initial Breast cancer Therapy Variation by Physician Specialty. *Am J Surg* 2013; 206(2): 218-222.
3. Jochelson M, Hayes DF, Ganz PA. Surveillance and Monitoring in Breast Cancer Survivors: Maximizing Benefit and Minimizing Harm. *ASCO Educational Book* 2013 e13 – e18.
4. Khatcheressian JL, Hurley P, Bantug E, et al.. Breast Cancer Follow-up and Management After Primary Treatment: American Society of Clinical Oncology Clinical Practice Guideline Update . *J Clin Oncol.* 2013 March 1; 31(7):961-965.
5. Bychkovsky BL, Lin NU. Imaging in the evaluation and follow-up of early and advanced breast cancer: When, why, and how often? *Breast.* 2017 Feb;31:318-324 Review.
6. Expert Panel on Breast Imaging: Moy L, Bailey L, D'Orsi C et al..ACR Appropriateness Criteria[®] Stage I Breast Cancer: Initial Workup and Surveillance for Local Recurrence and Distant Metastases in Asymptomatic Women. *J Am Coll Radiol.* 2017 May;14(5S):S282-S292.
7. Lam DL, Houssami N, Lee JM. Imaging Surveillance After Primary Breast Cancer Treatment. *AJR Am J Roentgenol.* 2017 Mar;208(3):676-686. Review.

Statement: Physical examination

1. Margenthaler JA, Allan E, Cheng L, et al.. Surveillance of Patients With Breast Cancer After Curative-Intent Primary Treatment: Current Practice Patterns. *Journal of Oncology Practice* 2012; 8(2): 79 – 83.
2. Khatcheressian JL, Hurley P, Bantug E, et al.. Breast Cancer Follow-up and Management After Primary Treatment: American Society of Clinical Oncology Clinical Practice Guideline Update . *J Clin Oncol.* 2013 March 1; 31(7):961-965.
3. Jochelson M, Hayes DF, Ganz PA. Surveillance and Monitoring in Breast Cancer Survivors: Maximizing Benefit and Minimizing Harm. *ASCO Educational Book* 2013 e13 – e18.
4. Parmeshwar R, Margenthaler JA, Allam E et al.. Patient Surveillance After Initial Breast cancer Therapy Variation by Physician Specialty. *Am J Surg* 2013; 206(2): 218-222.

Statement: Mammography

1. De Bock GH, Bonnema J, van Der Hage J et al.. Effectiveness of Routine Visits and Routine Tests in Detecting Isolated Locoregional Recurrences After Treatment for Early-Stage Invasive Breast Cancer: A Meta-Analysis and Systematic Review. *J Clin Oncol* 2004; 22(19): 4010-4018 .
2. Khatcheressian JL, Hurley P, Bantug E et al.:Breast Cancer Follow-up and Management After Primary Treatment: American Society of Clinical Oncology Clinical Practice Guideline Update . *J Clin Oncol.* 2013 March 1; 31(7):961-965.

Statement: Sonography of the breast

1. Graf O, Helbich TH, Fuchsjaeger MH et al.. Follow-up of palpable circumscribed noncalcified solid breast masses at mammography and US: can biopsy be averted? *Radiology* 2004; 233(3): 850-6.
2. Dillon MF, Hill AD, Quinn CM et al.: The accuracy of ultrasound, stereotactic, and clinical core biopsies in the diagnosis of breast cancer, with an analysis of false-negative cases. *Ann Surg.* 2005; 242(5):701-7.
3. Karella A, Vedantham S. Breast cancer imaging: a perspective for the next decade. *Med Phys* 2008; 35(11):4878-97. Review.
4. Jung Hyun-Yun., Min Jung Kim, Eun-Kyung Kim et al.. Imaging Surveillance of Patients with Breast Cancer after Primary Treatment: Current Recommendations. *Korean J Radiol* 2015;16(2):219-228.
5. Song SE, Cho N, Chang JM et al.. Diagnostic performances of supplemental breast ultrasound screening in women with personal history of breast cancer. *Acta Radiol.* 2017 Jan 1:284185117725779.

Statement: MRI of the breast

1. DeMartini W, Lehman C. A review of current evidence-based clinical applications for breast magnetic resonance imaging. *Top Magn*

- Reson Imaging 2008;19(3):143-50. Review.
2. Warner E. The role of magnetic resonance imaging in screening women at high risk of breast cancer. Top Magn Reson Imaging. 2008; 19(3):163-9. Review.
 3. Lehman CD, Lee JM, DeMartini WB et al. Screening MRI in Women With a Personal History of Breast Cancer. J Natl Cancer Inst. 2016 Jan 7;108(3).
 4. Shah C, Ahlawat S, Khan A et al.. The Role of MRI in the Follow-up of Women Undergoing Breast-conserving Therapy. Am J Clin Oncol. 2016 Jun;39(3):314-9.
 5. Cho N, Han W, Han BK et al.:Breast Cancer Screening With Mammography Plus Ultrasonography or Magnetic Resonance Imaging in Women 50 Years or Younger at Diagnosis and Treated With Breast Conservation Therapy. JAMA Oncol. 2017 Nov 1;3(11):1495-1502.
 6. Kim EJ, Kang BJ, Kim SH et al.:Diagnostic Performance of and Breast Tissue Changes at Early Breast MR Imaging Surveillance in Women after Breast Conservation Therapy.Radiology. 2017 Sep;284(3):656-666.
 7. Tadros A, Ardit B, Weltz C et al.:Utility of surveillance MRI in women with a personal history of breast cancer.Clin Imaging. 2017 Nov - Dec;46:33-36.

Statement: Pelvic examination Expert Opinion

1. Cohen I, Beyth Y, Tepper R.: The role of ultrasound in the detection of endometrial pathologies in asymptomatic postmenopausal breast cancer patients with tamoxifen treatment. Obstet Gynecol Surv 1998; 53(7): 429-38.
2. Giorda G, Crivellari D, Veronesi A et al.: Comparison of ultrasonography, hysteroscopy, and biopsy in the diagnosis of endometrial lesions in postmenopausal tamoxifen-treated patients. Acta Obstet Gynecol Scand 2002; 81(10):975-80.
3. Robertson C1, Arcot Ragupathy SK, Boachie C et al.: The clinical effectiveness and cost-effectiveness of different surveillance mammography regimens after the treatment for primary breast cancer: systemic reviews registry database analyses and economic evaluation. Health Technol Assess. 2011;15(34): 1-322.
4. Geurts SM, de Vegt F, Siesling S et al. Pattern of follow up care and early relapse detection in breast cancer patients. Breast Cancer Res Treat 2012; 136(3): 859-68.
5. Khatcheressian JL, Hurley P, Bantug E et al.. Breast Cancer Follow-up and Management After Primary Treatment: American Society of Clinical Oncology Clinical Practice Guideline Update . J Clin Oncol. 2013 March 1; 31(7):961-965.

Statement: Dexa scan Expert Opinion

1. Mahon SM, Williams MT, Spies MA: Screening for second cancers and osteoporosis in long-term survivors. Cancer Pract 2000; 8(6):

282-90.

2. Runowcz CD, Leach CR, Henry L et al.:American Cancer Society/American Society of Clinical Oncology breast cancer survivorship care guideline. CA Cancer J Clin 2016; 66: 43-73.
3. Shapiro CL, Van Poznak C, Lacchetti C, et al. Management of Osteoporosis in Survivors of Adult Cancers With Nonmetastatic Disease: ASCO Clinical Practice Guideline. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology* 2019;37:2916-46.



Routine-Nachsorgeuntersuchungen bei asymptomatischen Patientinnen

	Oxford	LoE	GR	AGO
▪ Routineleber (inkl. Tumormarker)	1a	A	-	-
▪ Labor zum Monitoring der Akut- und Spättoxizitäten der Therapien	5	D	+	+
▪ Lebersonographie	1a	A	-	-
▪ Skelettszintigraphie	1a	A	-	-
▪ Thorax-Röntgen	1a	A	-	-
▪ CT-Untersuchungen (Thorax, Abdomen und Becken)	2a	D	-	-
▪ Detektion isolierter / zirkulierender Tumorzellen	2a	D	-	-
▪ PET-CT	2b	B	-	-
▪ Ganzkörper-MRT	2b	B	-	-

1. Bychkovsky BL, Lin NU. Imaging in the evaluation and follow-up of early and advanced breast cancer: When, why, and how often? *Breast.* 2017 Feb;31:318-324. Review.
2. Lam DL, Houssami N, Lee JM. Imaging Surveillance After Primary Breast Cancer Treatment. *AJR Am J Roentgenol.* 2017 Mar;208(3):676-686. Review.
3. Expert Panel on Breast Imaging; Moy L, Bailey L, D'Orsi C, Green ED et al. ACR Appropriateness Criteria® Stage I Breast Cancer: Initial Workup and Surveillance for Local Recurrence and Distant Metastases in Asymptomatic Women. *J Am Coll Radiol.* 2017 May;14(5S):S282-S292.
4. Lafranconi A, Pylkkänen L, Deandrea S et al.. Intensive follow-up for women with breast cancer: review of clinical, economic and patient's preference domains through evidence to decision framework. *Health Qual Life Outcomes.* 2017 Oct 19;15(1):206.

Statement: Magnetic resonance imaging (MRI) of the breast

1. DeMartini W, Lehman C. A review of current evidence-based clinical applications for breast magnetic resonance imaging. *Top Magn Reson Imaging* 2008; 19(3):143-50. Review.
2. Warner E. The role of magnetic resonance imaging in screening women at high risk of breast cancer. *Top Magn Reson Imaging.* 2008; 19(3):163-9. Review.
3. Shah C, Ahlawat S, Khan A et al.. The Role of MRI in the Follow-up of Women Undergoing Breast-conserving Therapy. *Am J Clin Oncol.* 2016 Jun;39(3):314-9.

Statement: Routine biochemistry (incl. tumor markers)

1. McShane LM, Altman DG, Sauerbrei W et al..Statistics Subcommittee of the NCI-EORTC Working Group on Cancer Diagnostics. Reporting recommendations for tumor marker prognostic studies. *J Clin Oncol* 2005; 23(36): 9067-72.
2. Harris LN, Ismaila N, McShane LM et al..Use of Biomarkers to Guide Decisions on Adjuvant Systemic Therapy for Women With Early-stage Invasive Breast Cancer: American Society of Clinical Practice Guideline. *J Clin Oncol* 2016; 34(10): 1134-50.
3. Cardoso F, Kyriakides S, Ohno S, et al. Early breast cancer: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up^{&x2020;}. *Annals of Oncology* 2019;30:1194-220.

Statement: Ultrasound of the liver

Statement: Bone scan

Statement: Chest X-ray

Statement: CT of chest, abdomen and pelvis

1. Emens LA, Davidson NE. The follow-up of breast cancer. *Semin Oncol*. 2003; 30(3): 338-48. Review.
2. Moschetti I, Cinquini M, Lambertini M et al..Follow-up strategies for women treated for early breast cancer. *Cochrane Database Syst Rev* 2005 Jan 25;(1) Review. Update in: *Cochrane Database Syst Rev* 2016; (5) CD001768.
3. Dull B, Linkugel A, Margenthaler JA, Cyr AE. Overuse of Chest CT in Patients With Stage I and II Breast Cancer: An Opportunity to Increase Guidelines Compliance at an NCCN Member Institution. *J Natl Compr Canc Netw*. 2017 Jun;15(6):783-789.

Statement: Detection of isolated/circulating tumor cells

1. Janni W, Vogl FD, Wiedswang G et al..Persistence of disseminated tumor cells (DTC) in bone marrow (BM) during Follow-up predicts increased risk for relapse – Up-date of the pooled European data. *Clin Cancer Res* 2011; 17(9): 2967-76.
2. Rack B, Schindlbeck C, Jückstock J et al.. Circulating tumor cells predict survival in early average-to-high risk breast cancer patients. SUCCESS Study Group. *J Natl Cancer Inst*. 2014 May 15;106(5).

Statement: PET / WB-MRI

1. Ide M.Cancer screening with FDG-PET. *Q J Nucl Med Mol Imaging* 2006; 50(1): 23-7.
2. Schöder H, Gönen M. Screening for cancer with PET and PET/CT: potential and limitations. *J Nucl Med* 2007; 48 Suppl 1:4S-18S. Review.

3. Lei L, Wang X, Chen Z. PET/CT Imaging for Monitoring Recurrence and Evaluating Response to Treatment in Breast Cancer. *Adv Clin Exp Med.* 2016 Mar-Apr;25(2):377-82.
4. Cho IH, Kong EJ. Potential Clinical Applications of ¹⁸F-Fluorodeoxyglucose Positron Emission Tomography/Magnetic Resonance Mammography in Breast Cancer. *Nucl Med Mol Imaging.* 2017 Sep;51(3):217-226. Review.
5. Melsaether A, Moy L. Breast PET/MR Imaging. *Radiol Clin North Am.* 2017 May;55(3):579-589. Review.



AGO e. V.
in der DGGG e.V.
in der DKD e.V.
Guidelines Breast
Version 2021 10

www.ago-online.de
FÜR SICHERHEIT
LEBEN
FÜHREN

Background for toxicity management

Tamoxifen:	Cholesterin, Triglyceride, Bilirubin, ALAT, ASAT, gamma-GT
Aromatase-Inhibitors:	Cholesterin, Triglyceride, Bilirubin, ALAT, ASAT, gamma-GT
Anthracyclines:	pro-BNP, possibly Troponin
Trastuzumab:	pro-BNP, possibly Troponin
Checkpoint-Inhibitors:	Bilirubin, ALAT, ASAT, gamma-GT, Kreatinin, TSH, fT3/T4, Myoglobin



Früherkennung von potenziell heilbaren Erkrankungen

Oxford
LoE GR AGO

Lokoregionäre Rezidive (Thoraxwand, intramammäre Rezidive):

- | | | | |
|--|----|---|-----|
| • Inzidenz 7–20 %
(abhängig von der Zeit der Nachbeobachtung) | | | |
| • Brust-Selbst-Untersuchung | 5 | D | + |
| • Klin. Untersuchung, Mammographie & US | 1a | A | ++ |
| • Mamma-MR bei unklarer Mammographie / -sonographie* | 3a | B | +/- |

* Bei erhöhtem Risiko erwägen (Alter unter 50 J., HR+, Beurteilbarkeit in MG+US C/D)

Statement incidence

1. Perry NM. Quality assurance in the diagnosis of breast disease. EUSOMA Working Party. Eur J Cancer 2001; 37: 159-172
2. Wapnir IL, Anderson SJ, Mamounas EP et al.. Prognosis after ipsilateral breast tumor recurrence and locoregional recurrences in five National Surgical Adjuvant Breast and Bowel Project node-positive adjuvant breast cancer trials. J Clin Oncol 2006; 24:2028-2037

Statement breast self examination

1. Thomas DB, Gao DL, Ray RM et al.: Randomized trial of breast self-examination in Shanghai: final results. J Natl Cancer Inst 2002; 94(19): 1445-57.
2. Khatcheressian JL, Wolff AC, Smith TJ. American Society of Clinical Oncology 2006 update of the breast cancer follow-up and management guidelines in the adjuvant setting. J Clin Oncol. 2006 Nov 1;24(31):5091-7.
3. Montgomery DA, Krupa K, Cooke TG.: Follow-up in breast cancer: does routine clinical examination improve outcome? A systematic review of the literature. Br J Cancer 2007; 97(12): 1632-41.

Statement physical examination, mammography & US & MRI

1. Beinart G, Gonzalez-Angulo AM, Broglio K.: Clinical course of 771 patients with bilateral breast cancer: characteristics associated with overall and recurrence-free survival. Clin Breast Cancer 2007; 7(11): 867-74.

2. Montgomery DA, Krupa K, Cooke TG.: Follow-up in breast cancer: does routine clinical examination improve outcome? A systematic review of the literature. *Br J Cancer*. 2007; 97(12): 1632-41.
3. Buist DS1, Abraham LA, Barlow WE et al.:Diagnosis of second breast cancer events after initial diagnosis of early stage breast cancer. Breast Cancer Surveillance Consortium. *Breast Cancer Res Treat* 2010; 124(3): 863-73.
4. Kim JY, Cho N, Koo HR et al..Unilateral breast cancer: screening of contralateral breast by using preoperative MR imaging reduces incidence of metachronous cancer. *adiology*. 2013 Apr;267(1):57-66.
5. Khatcheressian JL, Hurley P, Bantug E et al.: Breast Cancer Follow-up and Management After Primary Treatment: American Society of Clinical Oncology Clinical Practice Guideline Update . *J Clin Oncol*. 2013 March 1; 31(7):961-965.
6. Neuman HB, Schumacher JR, Francescatti AB et al.:Utility of Clinical Breast Examinations in Detecting Local-Regional Breast Events After Breast-Sохранation in Women with a Personal History of High-Risk Breast Cancer. *Ann Surg Oncol*. 2016 Oct;23(10):3385-91.
7. Tsai WC, Wei HK, Hung CF et al.:Better Overall Survival for Breast Cancer Patients by Adding Breast Ultrasound to Follow-Up Examinations for Early Detection of Locoregional Recurrence-A Survival Impact Study. *Ultrasound Med Biol*. 2016 Sep;42(9):2058-64.
8. Freedman RA, Keating NL, Partridge AH et al.:Surveillance Mammography in Older Patients With Breast Cancer-Can We Ever Stop?: A Review. *JAMA Oncol*. 2017 Mar 1;3(3):402-409.
9. Taros A, Ardit B, Weltz C et al.:Utility of surveillance MRI in women with a personal history of breast cancer. *Clin Imaging*. 2017 Nov - Dec;46:33-36.



Früherkennung von potenziell heilbaren Erkrankungen

Oxford
LoE GR AGO

Kontralaterales Mammakarzinom:

<ul style="list-style-type: none"> ▪ Rel. Risiko: 2,5–5 ▪ Inzidenz: 0,5–1,0 % / Jahr ▪ Brust-Selbst-Untersuchung ▪ Klin. Untersuchung, Mammographie & US ▪ Mamma-MR* <p>Männliches Mammakarzinom: Vorgehen analog wie beim Karzinom der Frau bei unklarer Mammographie / -sonographie**</p> <p>* Bei erhöhtem Risiko erwägen (Alter unter 50 J., HR+, Beurteilbarkeit in MG+US C/D) ** S. Kapitel „Brustkrebs: Spezielle Situationen/Männliches Mammakarzinom“</p>	5 D + 1a A ++ 3b B +/- 5 D +
--	---

Statement risk and incidence

1. Hooning MJ, Aleman BM, Hauptmann M et al.: Roles of radiotherapy and chemotherapy in the development of contralateral breast cancer J Clin Oncol 2008; 26(34): 5561-8.
2. Yerushalmi R, Kennecke H, Woods R et al.: Does multicentric/multifocal breast cancer differ from unifocal breast cancer? An analysis of survival and contralateral breast cancer incidence. Breast Cancer Res Treat 2009; 117(2): 365-70.
3. Bertelsen L, Mellemkjær L, Christensen J et al.: Age-Specific Incidence of Breast Cancer in Breast Cancer Survivors and Their First-Degree Relatives. Epidemiology 2009; 20(2): 175 – 80.
4. Chao C, Bhatia S, Xu L, et al. Incidence, Risk Factors, and Mortality Associated With Second Malignant Neoplasms Among Survivors of Adolescent and Young Adult Cancer. JAMA Network Open 2019;2:e195536-e.

Statement breast self examination

1. Thomas DB, Gao DL, Ray RM et al.: Randomized trial of breast self-examination in Shanghai: final results. J Natl Cancer Inst 2002; 94(19): 1445-57.
2. Montgomery DA, Krupa K, Cooke TG et al.: Follow-up in breast cancer: does routine clinical examination improve outcome? A systematic review of the literature. Br J Cancer 2007; 97(12): 1632-41.
3. Khatcheressian JL, Hurley P, Bantug E et al.: Breast Cancer Follow-up and Management After Primary Treatment: American Society of Clinical Oncology Clinical Practice Guideline Update. J Clin Oncol. 2013 March 1; 31(7):961-965.

Statement physical examination, mammography & US&MRI

1. Beinart G, Gonzalez-Angulo AM, Broglio K et al.: Clinical course of 771 patients with bilateral breast cancer: characteristics associated with overall and recurrence-free survival. *Clin Breast Cancer* 2007; 7(11): 867-74.
2. Montgomery DA, Krupa K, Cooke TG et al.: Follow-up in breast cancer: does routine clinical examination improve outcome? A systematic review of the literature. *Br J Cancer*. 2007; 97(12): 1632-41.
3. Khatcheressian JL, Hurley P, Bantug E, et al.: Breast Cancer Follow-up and Management After Primary Treatment: American Society of Clinical Oncology Clinical Practice Guideline Update. *J Clin Oncol*. 2013 March 1; 31(7):961-965.
4. Kim JY1, Cho N, Koo HR et al. Unilateral breast cancer: screening of contralateral breast by using preoperative MR imaging reduces incidence of metachronous cancer. *Radiology*. 2013 Apr;267(1):57-66.
5. Freedman RA, Keating NL, Partridge AH et al.: Surveillance Mammography in Older Patients With Breast Cancer-Can We Ever Stop?: A Review. *JAMA Oncol*. 2017 Mar 1;3(3):402-409.
6. Vapiwala N, Hwang WT, Kushner CJ et al.: No impact of breast magnetic resonance imaging on 15-year outcomes in patients with ductal carcinoma in situ or early-stage invasive breast cancer managed with breast conservation therapy. *Cancer*. 2017 Apr 15;123(8):1324-1332.
7. van Bodegraven EA, van Raaij JC, Van Goethem M et al.: Guidelines and recommendations for MRI in breast cancer follow-up: A review. *Eur J Obstet Gynecol Reprod Biol*. 2017 Nov;218:5-11.

Statement surveillance of male breast cancer

1. Ferzoco RM, Ruddy KJ, Optimal delivery of male breast cancer follow-up care: improving outcomes, *Breast Cancer: Targets and Therapy* 2015;7 371–379
2. Auvinen A, Curtis RE, Ron E. Risk of subsequent cancer following breast cancer in men. *J Natl Cancer Inst*. 2002;94(17):1330–1332.



Früherkennung von potenziell heilbaren Erkrankungen

Oxford
LoE GR AGO

Sonstige Zweitkarzinome:

- | | | | |
|--|----|---|----|
| • MDS (RR 10,9), AML (RR 2,6–5,3), Kolorektal RR 3,0; Endometrium RR 1,6; Ovar RR 1,5; Lymphome RR 7 | | | |
| • Screening auf Zweitmalignome entsprechend der gültigen Leitlinien | 5 | D | ++ |
| • Gyn. Krebsfrüherkennungsuntersuchung | 5 | D | ++ |
| • Routinemäßige transvaginale Sonographie / Biopsie des Endometriums | 1b | B | - |

Statement: Risk

1. Brown LM, Chen BE, Pfeiffer RM et al.: Risk of second non-hematological malignancies among 376,825 breast cancer survivors. *Breast Cancer Res Treat* 2007; 106(3): 439-51.
2. Kirova YM, De Rycke Y, Gambotti L et al.: Second malignancies after breast cancer: the impact of different treatment modalities. *Br J Cancer* 2008 Mar 11; 98(5): 870-4.
3. Schaapveld M, Visser O, Louwman MJ et al.: Risk of new primary nonbreast cancers after breast cancer treatment: a Dutch population-based study. *J Clin Oncol* 2008; 26(8): 1239-46.
4. Andersson M, Jensen MB, Engholm G et al.: Risk of second primary cancer among patients with early operable breast cancer registered or randomised in Danish Breast Cancer cooperative Group (DBCG) protocols of the 77, 82 and 89 programmes during 1977-2001. *Acta Oncol* 2008; 47(4): 755-64.
5. CG Valentini, L Fianchi, MT Voso et al.: Incidence of Acute Myeloid Leukemia after Breast Cancer, *Mediterr J Hematol Infect Dis*. 2011; 3(1): e2011069.
6. Kaplan H1, Malmgren J, De Roos AJ.: Risk of myelodysplastic syndrome and acute myeloid leukemia post radiation treatment for breast cancer: a population-based study. *Breast Cancer Res Treat*. 2013 Feb;137(3):863-7.
7. Kaplan HG, Malmgren JA, Atwood MK.: Increased incidence of myelodysplastic syndrome and acute myeloid leukemia following breast cancer treatment with radiation alone or combined with chemotherapy: a registry cohort analysis 1990-2005. *BMC Cancer*. 2011 Jun 21;11:260. doi: 10.1186/1471-2407-11-260.
8. Freedman RA, Seisler DK, Foster JC et al.: Risk of acute myeloid leukemia and myelodysplastic syndrome among older women receiving anthracycline-based adjuvant chemotherapy for breast cancer on Modern Cooperative Group Trials (Alliance A151511).

- Breast Cancer Res Treat. 2017 Jan;161(2):363-373. doi: 10.1007/s10549-016-4051-1.
9. Sung H, Hyun N, Leach CR, Yabroff KR, Jemal A. Association of First Primary Cancer With Risk of Subsequent Primary Cancer Among Survivors of Adult-Onset Cancers in the United States. JAMA 2020;324:2521-35.

Statement: Screening for secondary malignancies according to current guidelines

1. Runowicz CD, Leach CR, Henry NL, et al. American Cancer Society/American Society of Clinical Oncology Breast Cancer Survivorship Care Guideline. Journal of Clinical Oncology 2016;34:611-35.

Statement: Pelvic examination and PAP smear

1. Gerber B, Krause A, Müller H et al.: Ultrasonographic detection of asymptomatic endometrial cancer in postmenopausal patients offers no prognostic advantage over symptomatic disease discovered by uterine bleeding. Eur J Cancer 2001; 37(1): 64-71.
2. Fishman DA, Cohen L, Blank SV et al.: The role of ultrasound evaluation in the detection of early-stage epithelial ovarian cancer. Am J Obstet Gynecol 2005; 192(4): 1214-21.
3. Rieck GC, Lim K, Rogers MT et al.: Screening for familial ovarian cancer--management and outcome of women with moderate to high risk of developing ovarian cancer. Int J Gynecol Cancer 2006;16 Suppl 1: 86-91.
4. Chan JK, Manuel MR, Cheung MK et al.: Breast cancer followed by corpus cancer: is there a higher risk for aggressive histologic subtypes? Gynecol Oncol 2006; 102(3): 508-12.

Statement: Endometrial ultrasound / biopsy

1. Gerber B, Krause A, Müller H, et al.: Effects of adjuvant tamoxifen on the endometrium in postmenopausal women with breast cancer: a prospective long-term study using transvaginal ultrasound. J Clin Oncol 2000; 18(20): 3464-70
2. Barakat RR, Gilewski TA, Almadrones L et al.: Effect of adjuvant tamoxifen on the endometrium in women with breast cancer: a prospective study using office endometrial biopsy. J Clin Oncol 2000;18(20): 3459-63.
3. Fung MF, Reid A, Faught W et al.: Prospective longitudinal study of ultrasound screening for endometrial abnormalities in women with breast cancer receiving tamoxifen. Gynecol Oncol 2003; 91(1): 154-9.

Statement: Marrow neoplasms after adjuvant breast cancer therapy

1. Wolff AC, Blackford AL, Visvanathan K et al.: Risk of marrow neoplasms after adjuvant breast cancer therapy: the national comprehensive cancer network experience. J Clin Oncol. 2015; 33(4): 340-8.

Statement: Secondary lung tumors:

1. Hoekstra N, Fleury E, Merino Lara TR et al.: Long-term risks of secondary cancer for various whole and partial breast irradiation techniques. *Radiother Oncol.* 2018 Sep;128(3):428-433
2. Burt LM, Ying J, Poppe MM, Suneja G, Gaffney DK. Risk of secondary malignancies after radiation therapy for breast cancer: Comprehensive results. *Breast (Edinburgh, Scotland)* 2017;35:122-9.



Nachsorge bei invasiven und nicht invasiven Karzinomen

Synopsis

Empfehlung für asymptomatische Patientinnen
(mod. nach ASCO-ACS Empfehlungen 2016, NCCN 2021, ESMO2019 und S3-Leitlinie 2017)

	Nachsorge/Follow-Up*			Screening/ Follow up	
Jahre nach Primärtherapie	1	2	3	4 – 5	> 5
Anamnese, klinische Untersuchung, Beratung	inv.; alle 3 Mon.	in situ: alle 6 Mon.		alle 6 Mon.	alle 12 Mon.
Selbstuntersuchung	monatlich				
Bildgebende Diagnostik, Laboruntersuchungen	indiziert nur bei Symptomatik +/- Befunden +/- Verdacht auf Rezidiv/Metastasen bzw. Monitoring der Nebenwirkungen der Therapie				
Mammographie und ergänzende Sonographie	BET**	beidseits: alle 12 Monate			
Mastektomie		kontralateral alle 12 Monate			
Echokardiografie	6–12 Monate bzw. 5 Jahre nach Abschluß der Anthrazyklin- oder Trastuzumab-haltiger Therapie (v.a. bei Risikofaktoren)				

* Fortlaufende "Nachsungsuntersuchungen" bei noch laufender adjuvante Therapie

** nach BET: Erste Mammographie 3 Jahr nach initialer Mammographie, oder zumindest 6 Monate nach abgeschlossener Radiatio

- Runowcz CD, Leach CR, Henry L et al.: American Cancer Society/American Society of Clinical Oncology breast cancer survivorship care guideline. CA Cancer J Clin 2016; 66: 43-73
- Merkow RP, Korenstein D, Yeahia R et al.: Quality of Cancer Surveillance Clinical Practice Guidelines: Specificity and Consistency of Recommendations. JAMA Intern Med. 2017 May 1;177(5):701-709.
- Muradali D, Kennedy EB, Eisen A et al.: Breast screening for survivors of breast cancer: A systematic review. Prev Med. 2017 Oct;103:70-75.
- NCCN Clinical Practice Guidelines in Oncology, Breast Cancer Version 15.01.2021
https://www.nccn.org/professionals/physician_gls/pdf/breast.pdf
- Cardoso F, Kyriakides S, Ohno S, et al. Early breast cancer: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. Annals of Oncology 2019;30:1194-220.
- Interdisziplinäre S3-Leitlinie für die Früherkennung, Diagnostik, Therapie und Nachsorge des Mammakarzinoms. Langversion 4.0 – Dezember 2017 AWMF-Registernummer: 032-045OL; http://www.leitlinienprogramm-onkologie.de/fileadmin/user_upload/LL_Mammakarzinom_Langversion_4.0.pdf



Brustkrebs Nachsorge Dauer und „Breast Nurses“

	Oxford		
	LoE	GR	AGO
▪ Dauer der Nachsorge			
• Bis zu 5 Jahre	1c	A	++
• Bis zu 10 Jahre	1c	A	+
▪ Nachsorge durch spezialisierte „Breast nurses“	2b	B	+/-*

* Studien empfohlen

1. Sheppard C. Breast cancer follow-up: literature review and discussion. Eur J Oncol Nurs 2007;11(4):340-7.
2. van Hezewijk M, Ranke GM, van Nes JG et al.: Patients' needs and preferences in routine follow-up for early breast cancer; an evaluation of the changing role of the nurse practitioner. Eur J Surg Oncol 2011; 37(9): 765-73.
3. Interdisziplinäre S3-Leitlinie für die Früherkennung, Diagnostik, Therapie und Nachsorge des Mammakarzinoms. Langversion 4.0 – Dezember 2017 AWMF-Registernummer: 032-045OL; http://www.leitlinienprogramm-onkologie.de/fileadmin/user_upload/LL_Mammakarzinom_Langversion_4.0.pdf
4. NCCN Clinical Practice Guidelines in Oncology, Breast Cancer Version 3.17-10.17; https://www.nccn.org/professionals/physician_gls/pdf/breast.pdf



Luminal-like, HER2-positive and Triple-negative Breast Cancer Patients

- Intrinsic typing of breast cancer leads to subgroups with different course of disease. Thus, postoperative surveillance should be adapted to specific time-dependent hazards of recurrence.
- ER-positive patients have stable risk over many years requiring long term surveillance.
- However, patients with HER2-positive disease and TNBC have more risk in the early phase of follow-up and should therefore receive more intense surveillance in the first years of follow-up.

Ribelles et al. BCR 2013

1. Voduc KD, Cheang MC, Tyldesley S et al.: Breast cancer subtypes and the risk of local and regional relapse. *J Clin Oncol* 2010; 28(10): 1684-91.
2. Ribelles N, Perez-Villa L, Jerez JM et al.: Pattern of recurrence of early breast cancer is different according to intrinsic subtype and proliferation index. *Breast Cancer Res* 2013;15(5):R98.
3. Metzger-Filho O1, Sun Z, Viale G et al.: Patterns of Recurrence and Outcome According to Breast Cancer Subtypes in Lymph Node-Negative Disease: Results From International Breast Cancer Study Group Trials VIII and IX. *J Clin Oncol* 2013; 31(25): 3083-3090.
4. Benveniste AP, Dryden MJ, Bedrosian I et al.: Surveillance of women with a personal history of breast cancer by tumour subtype. *Clin Radiol*. 2017 Mar;72(3):266.e1-266.e6.
5. Chu AJ, Chang JM, Cho NK et al.: Imaging Surveillance for Survivors of Breast Cancer: Correlation between Cancer Characteristics and Method of Detection. *J Breast Cancer*. 2017 Jun;20(2):192-197.