



Diagnosis and Treatment of Patients with early and advanced Breast Cancer

© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2022.1E

Options for Primary Prevention: Modifiable Lifestyle Factors

Prevention

© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2022.1E

- **Versions 2011–2021:**
**Dall / Diel / Gerber / Hanf / Maass / Mundhenke / Rhiem / Solbach /
Solomayer / Thomssen / von Minckwitz**
- **Version 2022:**
Dall / Gerber

Risk Factors for Breast Cancer 1

© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2022.1E

- Older age
 - Genetics
 - Family history of cancer
 - Personal history of breast lesions
 - Non-proliferative lesions
 - Proliferative lesions w/o atypia
 - High risk lesions (ADH, LIN)
 - Breast cancer (DCIS, Inv. BC)
 - Breast density
 - Chest irradiation
 - Type II Diabetes mellitus
 - Hyperthyreoidism
 - Lifetime number of menstrual cycles
 - Early menarche, late menopause
 - Maternal pregnancy factors (e.g. pre-eclampsia) (risk reduction), and low physical activity during pregnancy (risk increase)
- ### Social risk factors
- Lower number of births or no pregnancy
 - Advanced age at first full term delivery

Risk Factors for Breast Cancer 2

© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2022.1E

- **Short duration or absence of breast feeding**
- **Postmenopausal BMI < 18.5 and > 25 and especially > 40 (obesity)**
- **Food content**
- **Steroid hormone therapy**
 - Recent oral contraceptive use
 - Hormone therapy (estrogen / gestagen combination) in postmenopausal women
- **Alcohol intake**
- **Nicotine**
- **Light exposure at night (night shifts) *contradictory***
- **Low physical activity**
- **Endocrine disruptors in fetal and early childhood development (e.g. DES, bisphenol-A, DDT)**
- **Effect of carcinogenic substances / working materials**
- **Exposition to ionizing radiation**



Deodorant-use and risk

Breast Cancer and Deodorants/Antiperspirants: a Systematic Review.

Allam MF¹: Cent Eur J Public Health. 2016 Sep;24(3):245-247. doi: 10.21101/cejph.a4475.

© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2022.1E



So far there is no evidence of a correlation between aluminum containing deodorants and breast cancer risk

- All observational studies that evaluated the association between breast cancer risk and deodorants / antiperspirants use were reviewed. We have only identified two case-control studies, carried out between 2002 and 2006.
- There was no risk of antiperspirants use in the pooled risk (odds ratio 0.40, 95 % confidence interval 0.35-0.46).
- Our comprehensive search has identified an insufficient number of studies to conduct a quantitative review and obtain reliable results. Further prospective studies are strongly needed.

www.ago-online.de

FORSCHEN
LEHREN
HEILEN

High Proportion of Postmenopausal Breast Cancer Attributable to Lifestyle Factors

population attributable fractions (PAFs) of modifiable risk factors

Risk factors: obesity, physical inactivity, alcohol, low-fiber intake, smoking

Results: retrospective cohort study (Netherlands Cancer Registry)

2000: subpopulations of obese women, inactive women, alcohol drinkers, smokers etc.

2010: breast cancer incidence as compared to background incidence in these subgroups

25.7 % of postmenopausal breast cancer cases in the Netherlands in 2010 were attributable to lifestyle factors

8.8 % attributed to obesity
6.6 % attributed to alcohol
5.5 % attributed to physical inactivity
3.2 % attributed to low fiber intake
4.6 % attributed to smoking

Update 2019: Tamimi et al, 2016
USA: more than a third of
postmenopausal breast cancers are
preventable through changes in
modifiable risk factors

van Germert et al., Int J Cancer 2015; 152: 155-162

Pregnancy Related Factors

Oxford

LoE

GR

Prevention

- **Any full-term pregnancy**
- **High number of pregnancies**
- **First full-term pregnancy before age of 30 years**
- **Breast feeding (protective if total breast-feeding time exceeds 1.5-2 years)**
- **Lower birth weight of the first born (3000-3500 vs. > 4500g RR = 1.53)**
- **Lower length of pregnancy first born (26-31. WOP vs. 40-41. WOP; HR = 2.38, p = 0.03)**

2b

B

2b

B

2b

B

3a

B

2b

B

2b

B

No influence

- **Polycystic Ovarian Syndrome PCO**
- **Assisted reproduction**
- **Abortion**

3b

C

2b

B

2b

B

Medical Primary Prevention*

© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2022.1E

- **ASS**
- **COX2-Inhibitoren**
- **Bisphosphonates**
- **Vitamin D**
- **Statins**

Oxford		
LoE	GR	AGO
2a	B	+/-
2a	B	+/-
2b	B	+/-
2b	B	+/-
2b	B	-

* No approval, consider side effects

Medical Prevention

© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2022.1E

Kehm RD et al., Regular use of aspirin and other non-steroidal anti-inflammatory drugs and breast cancer risk for women at familial or Genetic risk: a cohort study, Breast Cancer Res. 2019 Apr. 18;21(1):52

Prospective multinational cohort study, n = 5606, healthy women questionnaire, regular intake of ASS, NSAID, COX2-inhibitors

Regular ASS-intake: HR 0.61, CI 0.33-1.14, breast cancer incidence

Regular COX2-inhibitors : HR 0.39, CI 0.15-0.97, breast cancer incidence other NSAIDs: n.s.

[independent of BRCA-status]

The risk of breast, ovarian and endometrial cancer in obese women submitted to bariatric surgery: a meta-analysis

B Ishihara, D Farah, M Fonseca and A Nazário, Surg Obes Relat Dis 2020;16(10):1596-1602

© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2022.1E

- **Meta-analysis, of a total of 150,537 patients in the bariatric surgery arm and 1,461,938 women in the control arm.**
- **The risk of breast cancer was reduced by 49 % [RR: 0.39 (95 % CI [0.31 to 0.56]); $I^2 = 90$ %; 7 studies).**
- **The risk of ovarian cancer was reduced by 53 % [RR: 0.47 (95 % CI [0.27 to 0.81]); $I^2 = 0$ %; 3 studies).**
- **The risk of endometrial cancer was reduced by 67 % [RR: 0.33 (95 % CI [0.21 to 0.51]); $I^2 = 88$ %; 7 studies).**

Association of Body Fat and Risk of Breast Cancer in Postmenopausal Women With Normal Body Mass Index: A Secondary Analysis of a Randomized Clinical Trial and Observational Study.

Iyengar NM et al.: JAMA Oncol. 2019 Feb 1;5(2):155-163



© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2022.1E

- **WHI substudy**
- **Among the 3460 women included in the analysis (mean [SD] age, 63.6 [7.6] years), multivariable-adjusted hazard ratios for the risk of invasive breast cancer were 1.89 (95 % CI, 1.21-2.95) for the highest quartile of whole-body fat and 1.88 (95 % CI, 1.18-2.98) for the highest quartile of trunk fat mass.**
- **The corresponding adjusted hazard ratios for ER-positive breast cancer were 2.21 (95 % CI, 1.23-3.67) and 1.98 (95 % CI, 1.18-3.31), respectively.**

www.ago-online.de

**FORSCHEN
LEHREN
HEILEN**

Prevention by Changing Lifestyle Factors: Diet

Oxford

- **Preference of a balanced diet***
- **Mediterranean Diet**
- **Dietary components**
 - **Olive oil (extra virgin olive oil), as part of mediterranean diet**
 - **Fat reduced food**
 - **Reduced consumption of red meat**
 - **Nuts / peanuts (> 10g/d) (peanut butter without effect)**
 - **Fiber containing food**
 - **Vitamin D substitution for prevention (MaCa HR1,02)**
 - **Vegetables / fruits ****
 - **Phytoestrogens / soy**
 - **Vegetarian / vegan diet (no significant risk reduction)**
 - **Coffee (no significant reduction)**
 - **Supplementation of vitamins, minerals, trace elements**

LoE	GR	AGO
2b	B	+
2a	B	+
2b	B	+
2a	B	+
2b	C	+
2b	B	+
2a	B	+
1b	B	+/-
2a	B	+/-
2a	B	+/-
2b	C	+/-
2a	B	+/-
2a	B	-

* As recommended by German Society of Nutrition (DGE)

** Recommended as a part of healthy nutrition

Vitamin D Supplements and Prevention of Cancer and Cardiovascular Disease

N Engl J Med. 2019 Jan 3;380(1):33-44. doi: 10.1056/NEJMoa1809944. Epub 2018 Nov 10.

© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2022.1E

Randomized, placebo-controlled trial, with a two-by-two factorial design, of vitamin D₃ (cholecalciferol) at a dose of 2000 IU per day and marine n-3 (also called omega-3) fatty acids at a dose of 1 g per day

Primary end points were invasive cancer of any type and major cardiovascular events

25,871 participants

median follow-up of 5.3 years

124 breast cancers (Vit D group) vs. 122 (placebo group) Hazard Ratio: 1,02



Prevention by Modifying Lifestyle Risk Factors: Alcohol

© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2022.1E

- Reduction of alcohol intake reduces risk of breast cancer (ideal < 10g/d, class II evidence)

Particularly for

- ER+ / PR+ tumors
- Invasive lobular tumors

Oxford		
LoE	GR	AGO
2a	B	+
2a	B	

Nature, Nurture and cancer risks: Genetic and nutritional contributions to cancer

Theodoratou, E.: Annu Rev Nutr. 2017 August 21; 37: 293–320.
doi:10.1146/annurev-nutr-071715-051004



© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2022.1E



No association was classified as convincing (class I). The association between alcohol intake and ER+ breast cancer was classified as highly suggestive (Class II)

based on a meta-analysis of 20 prospective studies (≥ 30 g/d of alcohol consumption versus non-drinkers

RR (95 % CI): 1.35 (1.23, 1.48, p-value = 5.2×10^{-10} , $I^2 = 26$ %,

$P_{\text{small effect bias}} = 0.184$, $P_{\text{excess significance bias}} = 4 \times 10^{-8}$)

www.ago-online.de

**FORSCHEN
LEHREN
HEILEN**

Prevention by Modifying Lifestyle Risk Factors: Smoking

Oxford

LoE	GR	AGO
2a	B	++

- **Never smoking reduces risk of breast cancer (~ 15-24 % reduction of lifetime risk)**
- **Young women smoking have a 60 % increased risk of BC, when smoking > 10 years before the first childbirth (vs. never smokers)**

© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2022.1E

Smoking and Risk of Breast Cancer in the Generations Study Cohort

Jones, M.E.: Breast Cancer Res. 2017 Nov 22;19(1):118. doi: 10.1186/s13058-017-0908-4.

© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2022.1E



102,927 women recruited 2003–2013

average of 7.7 years of follow-up

The HR (reference group was never smokers) was
1.14 (95 % CI 1.03–1.25; $P = 0.010$) for ever smokers,
1.24 (95 % CI 1.08–1.43; $P = 0.002$) for starting smoking at ages < 17 years
1.23 (1.07–1.41; $P = 0.004$) for starting smoking 1–4 years after menarche

Women with a family history of breast cancer (ever vs never smokers HR 1.35; 95 % CI 1.12–1.62; $P = 0.002$) had a significantly larger HR ... than women without (ever smoker vs never smoker HR 1.07; 95 % CI 0.96–1.20; $P = 0.22$).

Prevention by Modifying Lifestyle Risk Factors: Physical Activity

Oxford

LoE	GR	AGO
2a	B	++

- **Physical exercise**

(Metabolic equivalents to 3–5 hrs
moderate pace walking per week)

These effects also apply to *BRCA1/2* mutation carriers and for women with an increased family risk.



Recreational Physical Activity Is Associated with Reduced Breast Cancer Risk in Adult Women at High Risk for Breast Cancer: A Cohort Study of Women Selected for Familial and Genetic Risk.

Kehm RD et al.: Cancer Res. 2020 Jan 1;80(1):116-125. doi: 10.1158/0008-5472.CAN-19-1847. Epub 2019 Oct 2.

© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2022.1E

- **Prospective cohort study**
- **N = 15550, women with fam. Hx of breast cancer**
- **multiplicative interactions of physical activity with predicted absolute breast cancer familial risk based on pedigree data and with BRCA1 and BRCA2 mutation status**
- **Higher physical activity → 20 % reduction of breast cancer incidence**
- **(HR0.80, CI 0.68-0.93), independent of BRCA-status or pedigree risk**

www.ago-online.de

**FORSCHEN
LEHREN
HEILEN**

Prevention by Modifying Lifestyle Risk Factors: Hormone Therapy in Postmenopausal Women

© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2022.1E

- **Avoiding hormonal therapy in postmenopausal women**

- **Avoiding estrogen / progestin combinations**
- **Avoiding estrogens only
(no increased, possibly reduced breast cancer risk, but increased risk for endometrial cancer, if not hysterectomized)**

Oxford		
LoE	GR	AGO
1b	A	+
1b	A	+/-



Epigenome-wide association study for lifetime estrogen exposure identifies an epigenetic signature associated with breast cancer risk.

Johansson A et al.: Clin Epigenetics. 2019 Apr 30;11(1):66.

© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2022.1E

Epidemiological data from EPIC-Italy (n = 31,864)

Study: estimated lifetime estrogen exposure

**Method: epigenome-wide association study, blood DNA samples, N = 216 ,
and 440 healthy controls**

**Results: an estimated 5 % increase in breast cancer risk per 1-year longer ELEE
(OR = 1.05, 95 % CI 1.04-1.07, P = 3×10^{-12}) in EPIC-Italy.**

694 CpG sites were associated with ELEE (FDR Q < 0.05)

www.ago-online.de

**FORSCHEN
LEHREN
HEILEN**

Prevention of Hormones in Postmenopausal Patients

© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2022.1E

	N	MC-RR (95%CI)	Further information
WHI WHI: JAMA 2002, JAMA 2017	~ 27 000	1.3 (1,0-1,6)	1.3 (1.1-1,6) coronary events 1.4 (1,1-1,9) insults 2.1 (1,4-3,3) pulmonary embolism 2.1 (1,5-2,9) deep vein thrombosis
HERS Hulley S: JAMA 2002	I 2763 RCT, med. 4.1 yrs. II 2321 open-label, 2.7 yrs.	1.2 (0.95-1.5)	med. age 67 yrs. no secondary prevention side effects as comp. to WHI + cholecystectomy ↗
Million Women Beral V: Lancet 2003	1.084 110 ~ 50 % HRT 4.1 J. follow-up	1.66 (1.6-1.8)	EPC > E mode of applic. not relevant duration > 5 yrs. Tibolon RR 1.45 (1.2-1.7)
EPIC Int J Cancer 2010	1.153 747 person-years	1.4 (1.2-1.6) 1.8 (1.4-2.2)	E-Mono EPC > E
Metaanalyse Nelson HD: JAMA 2002	16 Studies	1.21-1.40	side effects as compared to WHI +

www.ago-online.de

Prevention of Hormones (EGC) in Postmenopausal Patients

© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2022.1E

	N	MC-RR (95% CI)	Further statements
CLEAR-study (NSW)	1236 BC cases	2.09 (1.57-2.78)	current user
Case-Control-Study, retrospect. Australia		1.03 (0.82-1.28)	past user
		2.62 (1.56-4.38)	E/P combination
		1.80 (1.21-2.68)	E only

Prevention by Modifying Lifestyle Risk Factors: Oral Contraception (OC)

Oxford

LoE

1a

- OC does not increase the risk of mortality from breast cancer
- Risk of breast cancer slightly increased, risk of ovarian, endometrial cancer is decreased

1a⁽⁻⁾

Risk Reduction for Ipsi- and Contralateral Breast Cancer

Rationale: Women with breast cancer have an increased risk for a second primary

Oxford

	LoE	GR	AGO
■ Tamoxifen*	1a	A	+
■ Aromatase inhibitors*	1a	A	+
■ Suppression of ovarian function* + Tamoxifen	1b	B	+

* Only proven for ER / PgR-positive primary sporadic BC

© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2022.1E