

Diagnosis and Treatment of Patients with early and advanced Breast Cancer



© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2022.1E

Endocrine based and targeted Therapy of Metastatic Breast Cancer

Endocrine Therapy of Metastatic Breast Cancer

- **Versions 2002–2021:**

Albert / Bischoff / Dall / Fasching / Fersis / Friedrich / Gerber / Huober / Janni / Jonat / Kaufmann / Kolberg-Liedtke / Loibl / Lüftner / Lück / von Minckwitz / Möbus / Müller / Mundhenke / Nitz / Schmidt / Schneeweiß / Schütz / Stickeler / Thill / Untch / Wöckel

- **Version 2022:**

Schmidt / Witzel

Endocrine Therapy of Metastatic Breast Cancer

© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2022.1E

Indication

Oxford LoE: 1a

GR: A

AGO: ++

Endocrine-based therapy is first line treatment in patients with metastatic breast cancer and positive (or unknown) hormone receptor (HR) status.

Exception: imminent organ failure

Caveat: HR may change during the course of disease.

Histology of recurrent site should be obtained whenever possible

Comparison ER / PR and HER2 Metastasis vs. Primary Tumor (n = 5.521)

Meta-analysis based on 39 (mostly retrospective) analyses, exclusively comparing primary tumor and metastasis (no lymph nodes):

Pooled discordance proportions were:

- 19,3% (95 % CI 1/4 15.8% to 23.4%) for ER
- 30,9% (95% CI 1/4 26.6% to 35.6%) for PR
- 10,3% (95% CI 1/4 7.8% to 13.6%) for HER2

Pooled proportions of tumors shifting from positive to negative

- 22.5% (95% CI = 16.4% to 30.0%) for ER
- 49.4% (95% CI = 40.5% to 58.2%) for PR
- 21.3% (95% CI = 14.3% to 30.5%) for HER2

Pooled proportions of tumors shifting from negative to positive

- 21.5% (95% CI = 18.1% to 25.5%) for ER
- 15.9% (95% CI = 11.3% to 22.0%) for PR
- 9.5% (95% CI = 7.4% to 12.1%) for HER2



Endocrine Therapy

General Considerations

- **Within all lines of treatment, treatment options should consider prior endocrine therapies, age and comorbidities as well as the respective approval status.**
- **Premenopausal patients treated with GnRH analogues or after ovariectomy can be treated like postmenopausal patients.**

© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2022.1E

www.ago-online.de

FORSCHEN
LEHREN
HEILEN

Metastatic Breast Cancer

Endocrine Resistance

© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2022.1E

Primary endocrine resistance:

- Relapse within 2 years of adjuvant endocrine treatment (ETx)
- Progressive disease within first 6 months of first-line ETx for MBC

Secondary (required) endocrine resistance:

- Relapse while on adjuvant ETx but after the first 2 years or a relapse within 12 months after completing adjuvant ETx
- PD \geq 6 months after initiation of ET for MBC

Endocrine Therapy in Premenopausal Patients with HER2-Negative Metastatic Breast Cancer

Oxford

- GnRHa + Fulvestrant + CDK4/6i
- GnRHa + AI + Ribociclib
- GnRHa + AI + Palbociclib / Abemaciclib
- GnRHa + Tamoxifen + Palbociclib / Abemaciclib
- GnRHa + Tamoxifen
- Tamoxifen
- GnRHa + AI (first + second line)
- GnRHa + Fulvestrant
- Aromataseinhibitors without OFS

LoE	GR	AGO
2b	B	++
1b	B	++
3b/5	C	+
2b	B	+/-
1a	A	+
2b	B	+/-
2b	B	+
1b	B	+
3	D	--

Endocrine Mono-Therapy in Postmenopausal Patients with HER2-Negative Metastatic Breast Cancer



© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2022.1E

- **Fulvestrant 500 mg**
- **Aromatase inhibitor***
- **Tamoxifen**
- **Fulvestrant 250 mg + Anastrozole**
- **Repeat prior treatments**

Oxford		
LoE	GR	AGO
1b	B	+
1a	A	+
1a	A	+
1b	B	+/-
5	D	+/-

* There is no evidence for superiority of a single aromatase inhibitor. As everolimus plus exemestane is indicated after AI treatment, a non-steroidal AI should be used in first line.

Endocrine-Based Treatment Options for Postmenopausal Patients with HER2-Negative Metastatic Breast Cancer

© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2022.1E

www.ago-online.de

FORSCHEN
LEHREN
HEILEN

	Oxford		
	LoE	GR	AGO
<ul style="list-style-type: none"> ▪ CDK4/6-Inhibitor (Abemaciclib, Palbociclib, Ribociclib) <ul style="list-style-type: none"> ▪ + non-steroidal AI ▪ + Fulvestrant 	1a	A	++
	1a	A	++
▪ Abemaciclib monotherapy	3	C	+/-
▪ Alpelisib + Fulvestrant (PIK3CA mutated)	1b	B	+
▪ Everolimus <ul style="list-style-type: none"> ▪ + Exemestane ▪ + Tamoxifen ▪ + Letrozole ▪ + Fulvestrant 	1b	A	+
	2b	B	+
	2b	B	+/-
	2b	B	+
▪ CDK4/6-Inhibitor beyond progression	3b	C	+/-
▪ CDK4/6-Inhibitor switch based on toxicity	5	D	+/-

CDK4/6 Inhibitors in First-line Studies

© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2022.1E

Study name	Paloma-2	Monarch-3	Monaleesa-2	Monaleesa-7
Treatment arms	Palbociclib / placebo with letrozole	Abemaciclib / placebo with nonsteroidal AI	Ribociclib / placebo with letrozole	Ribociclib / placebo with tamoxifen or non-steroidal aromatase inhibitor, all with goserelin
Patients	666	493	668	672
Menopausal status	postmenopausal	postmenopausal	postmenopausal	premenopausal
Progression-free survival (months, m)	27.6 vs. 14.5 m (+ 13.1 m) (HR 0.563)	28.2 vs. 14.8 m (+ 13.4 m) (HR 0.540)	25.3 vs. 16.0 m (+ 9.3 m) (HR 0.568)	23.8 vs. 13.0 m (+ 10.8 m) (HR 0.55)
Overall survival (months, m)	not reported	not reported	63.9 vs. 51.4 m (+ 12.5 m) (HR 0.76)	58.7 vs. 48.0 m (+ 10.7 m) (HR, 0.76)

Endocrine Therapy in Postmenopausal HER2-Negative Metastatic Breast Cancer in Combination with Bevacizumab

© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2022.1E

- **Maintenance bevacizumab plus endocrine therapy after remission with chemotherapy and bevacizumab**
- **Bevacizumab plus endocrine treatment as first line therapy for advanced disease**

Oxford		
LoE	GR	AGO
1b	B	+/-
1b	B	+/-

PARP Inhibitors in Patients with HER2-negative, gBRCA-Mutant, Metastatic Breast Cancer



© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2022.1E

- **Olaparib**

Oxford		
LoE	GR	AGO
1b	A	++

- **Talazoparib**

Oxford		
LoE	GR	AGO
1b	A	++

Diagnosis and Treatment of Patients with early and advanced Breast Cancer



© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2022.1E

HER2-Positive and HR-Positive Metastatic Breast Cancer

Endocrine Therapy in Postmenopausal HER2-Positive Metastatic Breast Cancer Patients

© AGO e. V.
in der DGGG e.V.
sowie
in der DKG e.V.

Guidelines Breast
Version 2022.1E

- **Anastrozole + trastuzumab**
- **Letrozole + trastuzumab**
- **Letrozole + lapatinib**
- **Fulvestrant + lapatinib**
- **Abemaciclib + fulvestrant + trastuzumab (after T-DM1)**
- **Aromatase inhibitors + trastuzumab / pertuzumab***

Oxford		
LoE	GR	AGO
1b	B	+/-
2b	B	+/-
1b	B	+/-
1b	B	+/-
2b	B	+/-
2b	B	+/-

Poor efficacy of endocrine therapy alone.

Consider induction chemotherapy + anti-HER2-therapy (followed by endocrine + anti-HER2-therapy as maintenance therapy)!

* Study participation recommended

Concomitant or Sequential Endocrine-Cytostatic Treatment

Oxford

LoE GR AGO

- **Concomitant endocrine-cytotoxic treatment**

- May increase response rate and progression free interval but not overall survival
 - May increase toxicity

1b **A** **-**

- **Endocrine maintenance therapy after chemotherapy +/- anti-HER2 therapy-induced response +/- anti HER2 therapy**

- Increases progression free interval

2b **B** **+**